


Dileptons in low-energy heavy-ion-collisions

Hendrik van Hees

Goethe University Frankfurt

October 12, 2021



Outline

1 Electromagnetic probes

- Electromagnetic radiation from hot/dense QCD matter
- Hadronic many-body theory

2 Bulk-medium evolution with transport and coarse graining

- coarse-graining in UrQMD

3 Dileptons in heavy-ion collisions

- Dielectrons (SIS/NA49/NA61/NA62)
- Dielectrons at FAIR/RHIC-BES

4 Signatures of the QCD-phase structure?

5 Conclusions and Outlook

Electromagnetic probes theory perspective

Electromagnetic probes in heavy-ion collisions

- γ, ℓ^\pm : no strong interactions
- reflect whole “history” of collision:
 - from pre-equilibrium phase
 - from thermalized medium
QGP and hot hadron gas
 - from VM decays after thermal freezeout





Fig. by A. Drees

Electromagnetic probes from thermal source

- retarded electromagnetic-current-correlation function

$$\Pi_{\text{em},i}^{\mu\nu} = i \int d^4x \exp(iq \cdot x) \Theta(x^0) \langle [j_{\text{em},i}^\mu(x), j_{\text{em},i}^\nu(0)] \rangle$$

- McLerran-Toimela formula [MT85, GK91]

$$q_0 \frac{dN_\gamma}{d^4x d^3\vec{q}} = -\frac{\alpha_{\text{em}}}{2\pi^2} g^{\mu\nu} \text{Im } \Pi_{\mu\nu}^{(\text{ret})}(q, u) \Big|_{q_0=|\vec{q}|} f_B(q \cdot u)$$
$$\frac{dN_{e^+e^-}}{d^4x d^4q} = -g^{\mu\nu} \frac{\alpha^2}{3q^2\pi^3} \text{Im } \Pi_{\mu\nu}^{(\text{ret})}(q, u) \Big|_{q^2=M_{e^+e^-}^2} f_B(q \cdot u)$$

- Lorentz covariant (dependent on four-velocity of fluid cell, u)
- $q \cdot u = E_{\text{cm}}$: Doppler blue shift of q_T spectra!
- to lowest order in α : $4\pi\alpha\Pi_{\mu\nu} \simeq \Sigma_{\mu\nu}^{(\gamma)}$
- vector-meson dominance model:

$$\Sigma_{\mu\nu}^{\gamma} = \text{G}_\rho$$

- $\ell^+\ell^-$ -inv.-mass spectra \Rightarrow in-med. spectral functions of vector mesons (ρ, ω, ϕ)!

Radiation from thermal QGP: $q\bar{q}$ annihilation

- General: McLerran-Toimela formula


$$\frac{dN_{l^+l^-}^{(\text{MT})}}{d^4x d^4q} = -\frac{\alpha^2}{3\pi^3} \frac{L(M^2)}{M^2} g_{\mu\nu} \text{Im} \sum_i \Pi_{\text{em},i}^{\mu\nu}(M, \vec{q}) f_B(q \cdot u)$$

- in QGP phase: $q\bar{q}$ annihilation
 - hard-thermal-loop improved em. current-current correlator

$$-i\Pi_{\text{em}, \text{QGP}} = \gamma^* \text{ (wavy line)} \rightarrow q \text{ (green loop with arrows)} \rightarrow \gamma^* \text{ (wavy line)}$$

Radiation from hadronic sources: ρ , ω , ϕ decays



- model assumption: vector-meson dominance




$$\begin{aligned}\frac{dN_{\rho \rightarrow l^+ l^-}^{(\text{MT})}}{d^4x d^4q} &= \frac{M}{q^0} \Gamma_{\rho \rightarrow l^+ l^-}(M) \frac{dN_\rho}{d^3\vec{x} d^4q} \\ &= -\frac{\alpha^2}{3\pi^3} \frac{L(M^2)}{M^2} \frac{m_\rho^4}{g_\rho^2} g_{\mu\nu} \text{Im } D_\rho^{\mu\nu}(M, \vec{q}) f_B \left(\frac{q \cdot u - 2\mu_\pi(t)}{T(t)} \right)\end{aligned}$$

- special case of McLerran-Toimela (MT) formula
- $M^2 = q^2$: invariant mass, M , of dilepton pair
- $L(M^2) = (1 + 2m_l^2/M^2) \sqrt{1 - 4m_l^2/M^2}$: dilepton phase-space factor
- $D_\rho^{\mu\nu}(M, \vec{q})$: (four-transverse part of) in-medium ρ propagator at given $T(t)$, $\mu_{\text{meson/baryon}}(t)$
- analogous for ω and ϕ

Transition form factors: “ ρ mesons” via VMD



- vector mesons have “vacuum spectral shapes”
- propagated as “on-shell particles” of finite lifetime and variable mass
- **Dalitz decay:** 1 particle \rightarrow 3 particles
- $V: \omega \rightarrow \pi + \gamma^* \rightarrow \pi + \ell^+ + \ell^-$
- $P, S: \pi, \eta \rightarrow \gamma + \gamma^* \rightarrow \gamma + \ell^+ + \ell^-$
- $R:$ Baryon resonances $\Delta, N^* \rightarrow N + V \rightarrow N + \gamma^* \rightarrow N + \ell^+ + \ell^-$
- vector-meson dominance: model for hadron em. trans. FF




GiBUU: “ ρ meson” in pp

- production through hadron resonances

$$NN \rightarrow NR \rightarrow NN\rho, NN \rightarrow N\Delta \rightarrow NN\pi\rho$$


$\rho \rightarrow e^+e^-$



- plots: J. Weil et al [WHM12, ABB⁺14]
- VMD model \Leftrightarrow em. transition form factors of baryon resonances!
- “ ρ ”-line shape “modified” already in elementary hadronic reactions
- due to production mechanism via resonances


Hadronic many-body theory

- hadronic many-body theory (HMBT) for vector mesons
[Ko et al, Chanfray et al, Herrmann et al, Rapp et al, ...]
- $\pi\pi$ interactions and **baryonic excitations**
- effective hadronic models, implementing symmetries
- parameters fixed from phenomenology (photon absorption at nucleons and nuclei, $\pi N \rightarrow \rho N$)
- evaluated at **finite temperature and density**
- self-energies \Rightarrow **mass shift and broadening** in the medium





- Baryons** important, even at low **net** baryon density $n_B - n_{\bar{B}}$
- reason: $n_B + n_{\bar{B}}$ relevant (CP inv. of strong interactions)

Meson contributions



[RG99]

In-medium spectral functions and baryon effects




[RW99]

- **baryon effects** important
 - large contribution to broadening of the peak
 - responsible for most of the strength at small M

Dilepton rates: Hadron gas \leftrightarrow QGP

- in-medium **hadron gas** matches with **QGP**
- similar results also for γ rates
- “quark-hadron duality”?



[Rap13]

Bulk-medium evolution

Bulk evolution with transport and coarse graining

- established transport models for **bulk evolution**
 - e.g., UrQMD, GiBUU, BAMPS, (p)HSD,...
 - solve **Boltzmann equation** for hadrons and/or partons
- dilemma: need medium-modified **dilepton/photon emission rates**
- usually available only in **equilibrium QFT calculations**
- ways out:
 - **(ideal) hydrodynamics** \Rightarrow local thermal equilibrium \Rightarrow use equilibrium rates
 - transport-hydro hybrid model: treat early stage with transport, then **coarse grain** \Rightarrow switch to hydro \Rightarrow switch back to transport (**Cooper-Frye “particilization”**)
- here: **UrQMD transport** for entire bulk evolution
 \Rightarrow use **coarse graining** in space-time cells \Rightarrow extract T, μ_B, μ_π, \dots \Rightarrow use equilibrium rates locally

Coarse-grained UrQMD (CGUrQMD)


- problem with **medium modifications** of spectral functions/interactions
- only available in equilibrium many-body QFT models
- use “in-medium cross sections” naively: **double counting?!**
- way out: map transport to **local-equilibrium fluid**
- use **ensemble of UrQMD** runs with an **equation of state**
- space-time grid with $\Delta t = 0.2 \text{ fm}/c$, $\Delta x = 0.8 \text{ fm}$
- fit **temperature, chemical potentials, flow-velocity field** from anisotropic energy-momentum tensor [FMR_S13]

$$T^{\mu\nu} = (\epsilon + P_{\perp}) u^{\mu} u^{\nu} - P_{\perp} g^{\mu\nu} - (P_{\parallel} - P_{\perp}) V^{\mu} V^{\nu}$$

- thermal rates from **partonic/hadronic QFT** become applicable
- here: **extrapolated lattice QGP** and **Rapp-Wambach HMBT**
- caveat: **consistency between EoS, matter content of QFT model/UrQMD!**


Coarse-grained UrQMD (CGUrQMD)

- $T_c = 170$ MeV; $T > T_c \Rightarrow$ lattice EoS; $T < T_c \Rightarrow$ HRG EoS



Coarse-grained UrQMD (CGUrQMD)


- pressure anisotropy (for In+In @ SPS; NA60)



Dielectrons (SIS/HADES)


CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- coarse-graining method works at low energies!
- UrQMD-medium evolution + RW-QFT rates




CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- dielectron spectra from $\text{Ar} + \text{KCl}(1.76 \text{ AGeV}) \rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra
- $M_{ee} < 0.13 \text{ GeV}$




CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- dielectron spectra from $\text{Ar} + \text{KCl}(1.76 \text{ AGeV}) \rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra
- $0.13 \text{ GeV} M_{ee} < 0.3 \text{ GeV}$




CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- dielectron spectra from $\text{Ar} + \text{KCl}(1.76 \text{ AGeV}) \rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra
- $0.3 \text{ GeV} M_{ee} < 0.45 \text{ GeV}$




CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- dielectron spectra from $\text{Ar} + \text{KCl}(1.76 \text{ AGeV}) \rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra
- $0.45 \text{ GeV} M_{ee} < 0.65 \text{ GeV}$




CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- dielectron spectra from $\text{Ar} + \text{KCl}(1.76 \text{ AGeV}) \rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra
- $M_{ee} > 0.65 \text{ GeV}$




CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- dielectron spectra from $\text{Ar} + \text{KCl}(1.76 \text{ AGeV}) \rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra
- rapidity spectrum ($M_{ee} < 0.13 \text{ GeV}$)



CGUrQMD: Au+Au (1.23 AGeV) (SIS/HADES)




[SGRS18]

- good agreement between models and data
- consistency between two independent coarse-grained-UrQMD simulations
- based on same Rapp-Wambach in-medium rates

Dielectrons at RHIC-BES/FAIR/NICA



CGUrQMD: Au+Au ($E_{\text{lab}} = 2-35 \text{ AGeV}$)



NB: also photon spectra [\[EHB16b\]](#)

Outlook: Dilepton polarization

- $\lambda = (\Pi_T - \Pi_L) / (\Pi_T + \Pi_L)$
- for NA60 with fireball blast-wave parametrization




Signatures of the QCD-phase structure?

QCD phase structure from em. probes?

- hadronic observables like p_T spectra: “snapshot” of the stage after **kinetic freezeout**
- particle abundancies: **chemical freezeout**
- em. probes: emitted during the whole medium evolution
life time of the medium \Rightarrow “four-volume of the fireball”
- use CGUrQMD to study **system-size dependence**
- study AA collisions for different A [EHWB15]
- **“excitation functions”:**
systematics of $\ell^+\ell^-$ (and γ) emission vs. beam energy [EHB16b, RH16]
similar study in [GHR⁺16]
- **caveat:** phase transition not really implemented!!!

Scaling behavior of thermal-dilepton yield


- central collisions from C+C to Au+Au at $E_{\text{kin}} = 1.76 \text{ AGeV}$



- thermal-dilepton yield roughly $\propto V_{\text{therm}}^{(4)} \propto A^{4/3} \propto At_{\text{therm}} \propto N_{\pi^0}^{4/3}$
- at low(est) beam energies: lifetime of “medium” \cong time nuclei pass through each other

Mass-temperature relation in dilepton emission


- interplay between increasing volume and decreasing temperature of fireball
- in IMR ($T < m_\phi < M_{\ell^+\ell^-} < m_{J/\psi}$) biased towards **early hot stages**
- only “background”: correlated $D\bar{D}$ decays, some Drell-Yan
- otherwise emission from **thermal** QGP and hadronic sources
- invariant-mass slope \Leftrightarrow true **invariant** space-time averaged **temperature**
- no blueshift due to radial flow as in p_t spectra (e.g., photons)



[EHB16b, EHB16a]


Dilepton systematics in the beam-energy scan

- thermal-fireball model [RH16, EHB16a]
- invariant-mass slope in IMR \Rightarrow true temperature!
- no blue shift from radial flow as in p_T/m_T spectra



Dilepton systematics in the beam-energy scan

- thermal-fireball model [RH16]
- beam-energy scan at RHIC and lower energies at FAIR and
- dilepton yield as **fireball-lifetime clock**



Conclusions and Outlook

- General ideas

- em. probes \Leftrightarrow in-medium em. current-correlation function
- dual rates around T_c (compatible with χ symmetry restoration)
- medium modifications of ρ, ω, ϕ
- importance of baryon-resonance interactions

- Application to dileptons in HICs

- coarse-grained transport (here: CGUrQMD)
- allows use of thermal-QFT spectral VM functions
- applicable also at low collision energies
- allows use of thermal-QFT models for dilepton rates
- successful description from SIS to RHIC energies
- consistent description of M and m_T spectra!
- effective slope of M spectra ($1.5 \text{ GeV} < M < M_{J/\psi}$) provides $\langle T \rangle$
- beam-energy scan at RHIC and FAIR \Rightarrow signature of phase transition?

- Outlook

- signature of cross-over vs. 1st order (or even critical endpoint)???
- challenge: phase transition in (coarse-grained) transport???

Bibliography I

- [ABB⁺14] G. Agakishiev, et al., Baryon resonance production and dielectron decays in proton-proton collisions at 3.5 GeV, *Eur. Phys. J. A* **50** (2014) 82.
<https://doi.org/10.1140/epja/i2014-14082-1>
- [EHB16a] S. Endres, H. van Hees, M. Bleicher, Energy, centrality and momentum dependence of dielectron production at collider energies in a coarse-grained transport approach, *Phys. Rev. C* **94** (2016) 024912.
<https://doi.org/10.1103/PhysRevC.94.024912>
- [EHB16b] S. Endres, H. van Hees, M. Bleicher, Photon and dilepton production at the Facility for Proton and Anti-Proton Research and beam-energy scan at the Relativistic Heavy-Ion Collider using coarse-grained microscopic transport simulations, *Phys. Rev. C* **93** (2016) 054901.
<https://doi.org/10.1103/PhysRevC.93.054901>

Bibliography II

- [EHWB15] S. Endres, H. van Hees, J. Weil, M. Bleicher, Dilepton production and reaction dynamics in heavy-ion collisions at SIS energies from coarse-grained transport simulations, Phys. Rev. C **92** (2015) 014911.
<https://doi.org/10.1103/PhysRevC.92.014911>
- [FMRS13] W. Florkowski, M. Martinez, R. Ryblewski, M. Strickland, Anisotropic hydrodynamics, Nucl. Phys. A **904-905** (2013) 803c.
<https://doi.org/10.1016/j.nuclphysa.2013.02.138>
- [GHR⁺16] T. Galatyuk, P. M. Hohler, R. Rapp, F. Seck, J. Stroth, Thermal Dileptons from Coarse-Grained Transport as Fireball Probes at SIS Energies, Eur. Phys. J. A **52** (2016) 131.
<https://doi.org/10.1140/epja/i2016-16131-1>
- [GK91] C. Gale, J. I. Kapusta, Vector dominance model at finite temperature, Nucl. Phys. B **357** (1991) 65.
[https://doi.org/10.1016/0550-3213\(91\)90459-B](https://doi.org/10.1016/0550-3213(91)90459-B)

Bibliography III

- [MT85] L. D. McLerran, T. Toimela, Photon and Dilepton Emission from the Quark-Gluon Plasma: Some General Considerations, Phys. Rev. D **31** (1985) 545.
<https://doi.org/10.1103/PhysRevD.31.545>
- [Rap13] R. Rapp, Dilepton Spectroscopy of QCD Matter at Collider Energies, Adv. High Energy Phys. **2013** (2013) 148253.
<https://doi.org/10.1155/2013/148253>
- [RG99] R. Rapp, C. Gale, ρ properties in a hot meson gas, Phys. Rev. C **60** (1999) 024903.
<https://doi.org/10.1103/PhysRevC.60.024903>
- [RH16] R. Rapp, H. van Hees, Thermal Dileptons as Fireball Thermometer and Chronometer, Phys. Lett. B **753** (2016) 586.
<https://doi.org/10.1016/j.physletb.2015.12.065>
- [RW99] R. Rapp, J. Wambach, Low mass dileptons at the CERN-SPS: Evidence for chiral restoration?, Eur. Phys. J. A **6** (1999) 415.
<https://doi.org/10.1007/s100500050364>

Bibliography IV

- [SGRS18] F. Seck, T. Galatyuk, R. Rapp, J. Stroth, Thermal dileptons as QCD matter probes at SIS, *J. Phys. Conf. Ser.* **1024** (2018) 012011.
<https://doi.org/10.1088/1742-6596/1024/1/012011>
- [WHM12] J. Weil, H. van Hees, U. Mosel, Dilepton production in proton-induced reactions at SIS energies with the GiBUU transport model, *Eur. Phys. J. A* **48** (2012) 111.
<https://doi.org/10.1140/epja/i2012-12111-9>