

```

>
#
#####
# Maple-Worksheet zu Uebungsblatt 8, Aufgabe 2 (1/r^2-
Potential)
#
#####

restart;

A := L^2/(2*m) - alpha;

# c, (a)
s := sqrt(E/A) * sin((sqrt(2*m*A)/L)*phi + Pi/2);
plot(subs(L=1.0, m=1.0, alpha=0.25, E=1.0, phi0=0.0, 1/s), phi=
-2.0..+2.0, coords=polar,
view=[-2.0..2.0, -2.0..2.0]);


# c, (b)
s := sqrt(-E/A) * sinh((sqrt(-2*m*A)/L)*phi);
plot(subs(L=1.0, m=1.0, alpha=1.0, E=1.0, phi0=0.0, 1/s), phi=
+0.25..+10.0, coords=polar,
view=[-0.2..+1.0, -0.2..+1.0]);

# c, (c)
s := sqrt(E/A) * cosh((sqrt(-2*m*A)/L)*phi);
plot(subs(L=1.0, m=1.0, alpha=1.0, E=-1.0, phi0=0.0, 1/s), phi=
-10.0..+10.0, coords=polar,
view=[-0.2..+0.8, -0.5..+0.5]);


```

$$A := \frac{1}{2} \frac{L^2}{m} - \alpha$$

$$s := \sqrt{\frac{E}{\frac{1}{2} \frac{L^2}{m} - \alpha}} \cos\left(\frac{\sqrt{2} \sqrt{m \left(\frac{1}{2} \frac{L^2}{m} - \alpha\right)} \phi}{L}\right)$$



$$s := \sqrt{-\frac{E}{\frac{1}{2} \frac{L^2}{m} - \alpha}} \sinh\left(\frac{\sqrt{-2 m \left(\frac{1}{2} \frac{L^2}{m} - \alpha\right)} \phi}{L}\right)$$



$$s := \sqrt{\frac{E}{\frac{1}{2} \frac{L^2}{m} - \alpha}} \cosh\left(\frac{\sqrt{-2 m \left(\frac{1}{2} \frac{L^2}{m} - \alpha\right)} \phi}{L}\right)$$

