

The Secret of Mass: Can we Evaporate the Vacuum at RHIC?

Hendrik van Hees

Texas A&M University

February 24, 2007



Outline

The Beauty of Nature: Symmetries


Elementary Particles

The Fundamental Interactions

Heavy-Ion Collisions and the Quark-Gluon Plasma

The Beauty of Nature: Symmetries


- ▶ What is a **symmetry**?
- ▶ Geometry: Certain operations like rotations or translations, **do not change** a figure \Rightarrow then we say “it’s **symmetric**”!



- ▶ rotating the hexagon by an angle of 60° **doesn't change it** \Rightarrow **Symmetry!**

The Beauty of Nature: Symmetries


- ▶ What is a **symmetry**?
- ▶ Geometry: Certain operations like rotations or translations, **do not change** a figure \Rightarrow then we say “it’s **symmetric**”!



- ▶ rotating the hexagon by an angle of 60° **doesn't change it** \Rightarrow **Symmetry!**

The Beauty of Nature: Symmetries


- ▶ What is a **symmetry**?
- ▶ Geometry: Certain operations like rotations or translations, **do not change** a figure \Rightarrow then we say “it’s **symmetric**”!



- ▶ rotating the hexagon by an angle of 60° **doesn't change it** \Rightarrow **Symmetry!**

The Beauty of Nature: Symmetries


- ▶ What is a **symmetry**?
- ▶ Geometry: Certain operations like rotations or translations, **do not change** a figure \Rightarrow then we say “it’s **symmetric**”!



- ▶ rotating the hexagon by an angle of 60° **doesn't change it** \Rightarrow **Symmetry!**

The Beauty of Nature: Symmetries


- ▶ What is a **symmetry**?
- ▶ Geometry: Certain operations like rotations or translations, **do not change** a figure \Rightarrow then we say “it’s **symmetric**”!



- ▶ rotating the hexagon by an angle of 60° **doesn't change it** \Rightarrow **Symmetry!**

The Beauty of Nature: Symmetries


- ▶ What is a **symmetry**?
- ▶ Geometry: Certain operations like rotations or translations, **do not change** a figure \Rightarrow then we say “it’s **symmetric**”!



- ▶ rotating the hexagon by an angle of 60° **doesn't change it** \Rightarrow **Symmetry!**

The Beauty of Nature: Symmetries

- ▶ What is a **symmetry**?
- ▶ Geometry: Certain operations like rotations or translations, **do not change** a figure \Rightarrow then we say “it’s **symmetric**”!



- ▶ rotating the hexagon by an angle of 60° **doesn't change it** \Rightarrow **Symmetry!**

Symmetry of Natural Laws and Conservation Laws

- ▶ Mathematician **Emmy Noether** found an important relation between **symmetries and conservation laws**
 - ▶ Equation, describing the **behavior of an object in time**, does not change under an operation (**symmetry**) \Leftrightarrow a certain quantity stays **constant in time** (**conserved quantity**)

Symmetry of Natural Laws and Conservation Laws

- ▶ Mathematician **Emmy Noether** found an important relation between **symmetries and conservation laws**
 - ▶ Equation, describing the **behavior of an object in time**, does not change under an operation (**symmetry**) \Leftrightarrow a certain quantity stays **constant in time** (**conserved quantity**)
 - ▶ it works also the other way:
conserved quantity \Rightarrow **equations obey a symmetry**

Symmetry of Natural Laws and Conservation Laws

- ▶ Mathematician **Emmy Noether** found an important relation between **symmetries and conservation laws**
 - ▶ Equation, describing the **behavior of an object in time**, does not change under an operation (**symmetry**) \Leftrightarrow a certain quantity stays **constant in time** (**conserved quantity**)
 - ▶ it works also the other way:
conserved quantity \Rightarrow **equations obey a symmetry**
- ▶ example 1: Natural Laws **do not change with time** (equations look the same at any time) \Rightarrow **Conservation of Energy**

Symmetry of Natural Laws and Conservation Laws

- ▶ Mathematician **Emmy Noether** found an important relation between **symmetries and conservation laws**
 - ▶ Equation, describing the **behavior of an object in time**, does not change under an operation (**symmetry**) \Leftrightarrow a certain quantity stays **constant in time** (**conserved quantity**)
 - ▶ it works also the other way:
conserved quantity \Rightarrow **equations obey a symmetry**
- ▶ example 1: Natural Laws **do not change with time** (equations look the same at any time) \Rightarrow **Conservation of Energy**
- ▶ example 2: Natural Laws **do not change with position** (equations of motion look the same at any place) \Rightarrow **Conservation of momentum**

Spontaneous Symmetry Breaking



Emmy Noether

- ▶ Equations are **symmetric**, but not the **state of lowest energy**
- ▶ Conservation law still true, but **symmetry not realized**

Spontaneous Symmetry Breaking



Emmy Noether

- ▶ Equations are **symmetric**, but not the **state of lowest energy**
- ▶ Conservation law still true, but **symmetry not realized**
- ▶ example: **rotating** a piece of iron, no change of laws for atoms
- ▶ it can be a **magnet** \Rightarrow **specifies a direction**

Spontaneous Symmetry Breaking




Emmy Noether

- ▶ Equations are **symmetric**, but not the **state of lowest energy**
- ▶ Conservation law still true, but **symmetry not realized**
- ▶ example: **rotating** a piece of iron, no change of laws for atoms
- ▶ it can be a **magnet** \Rightarrow **specifies a direction**
- ▶ Heating the magnet, at a certain “critical temperature” **iron becomes suddenly unmagnetic**
- ▶ Phase transition \Rightarrow **Symmetry restored**

Ferromagnets


- ▶ Magnetizing Ferromagnet: Applying a magnetic field



- ▶ Little “elementary magnets” inside the magnet stay lined up even after switching off the magnetic field
- ▶ Ferromagnet itself becomes a magnet
- ▶ specifies direction in space \Rightarrow spontaneous breaking of rotational symmetry

Ferromagnets


- ▶ heating up the ferro magnet rattles up the elementary magnets



- ▶ at a certain **critical temperature**: Alignment of elementary magnets lost (**phase transition!**)
- ▶ Iron rod is **no longer a magnet!**
- ▶ no direction specified anymore
- ▶ **rotational symmetry restored**

Elementary particles

- ▶ Since ancient times scientists have asked:
Are there indivisible smallest lumps of matter?
- ▶ Democritus (460-370 BC):
“There is nothing but atoms and empty space (the void).”
- ▶ atom=Greek for indivisible
- ▶ Rutherford (1909-1911):
most of the atom is “empty space”
- ▶ **mass** concentrated in the atomic nucleus



Subatomic particles


- ▶ **electrons are elementary**
- ▶ atomic nucleus is composed of **nucleons=protons and neutrons**
- ▶ nucleons made of **up** and **down** quarks



proton = (uud)



neutron = (udd)



- ▶ **up** quark: charge $+2/3$, mass $m_u = 3 \text{ MeV}/c^2$
- down** quark: charge $-1/3$, mass $m_d = 6 \text{ MeV}/c^2$
- electron: charge -1 , mass $m_e = 0.5 \text{ MeV}/c^2$

Subatomic particles


- ▶ **electrons are elementary**
- ▶ atomic nucleus is composed of **nucleons=protons and neutrons**
- ▶ nucleons made of **up** and **down** quarks



proton = (uud)




neutron = (udd)



- ▶ **up** quark: charge $+2/3$, mass $m_u = 3 \text{ MeV}/c^2$
- ▶ **down** quark: charge $-1/3$, mass $m_d = 6 \text{ MeV}/c^2$
- ▶ electron: charge -1 , mass $m_e = 0.5 \text{ MeV}/c^2$
- ▶ **BUT:** nucleon mass $m_p = m_n = 940 \text{ MeV}/c^2$


Elementary Particles and Fundamental Interactions

- ▶ What holds the particles together (forming **matter**)?
- ▶ Fundamental forces or interactions (see Professor Fries's talk!)
- ▶ Laws ruled by symmetries!
- ▶ e.g. **electric charge conserved** \Leftrightarrow "Force Carrier" (wave fields \leftrightarrow particles) for electromagnetic interaction **Photon**



	Gravity	Weak (Electroweak)	Electromagnetic	Strong
Carried By	Graviton (not yet observed)	$W^+ W^- Z^0$	Photon	Gluon
Acts on	All	Quarks and Leptons	Quarks and Charged Leptons and $W^+ W^-$	Quarks and Gluons

Matter particles vs. Force Carriers



- ▶ Elementary Matter Particles:
 - Quarks and Leptons
 - spin $s = 1/2$
- ▶ Elementary Force particles
 - gluons, photons (γ), W, Z
 - Spin $s = 1$

Matter particles vs. Force Carriers

ELEMENTARY PARTICLES				
Quarks	u	c	t	Force Carriers
	d	s	b	
Leptons	ν_e	ν_μ	ν_τ	Z
	e	μ	τ	W



- ▶ Elementary Matter Particles:
Quarks and Leptons
spin $s = 1/2$
- ▶ Elementary Force particles
gluons, photons (γ), W, Z
Spin $s = 1$
- ▶ Fermions: only one identical fermion per room!
Space-time symmetries:
particles with half-integer spin
- ▶ Bosons: identical bosons prefer to stay together!
Space-time symmetries:
particles with integer spin



The Eightfold Way

- ▶ in the 1950-1960'ies a **whole zoo of particles** has been discovered using accelerators (see Prof. Cagliardi's Talk!)
- ▶ most of them: **hadrons: particles participating in strong interaction**
- ▶ Gell-Mann, Zweig, Ne'eman (1961): all the hadrons can be understood by assuming that they are composed of **spin-1/2 particles with electric charges $-1/3$ and $2/3$**
- ▶ Gell-Mann: How to name them? **Quarks!**
- ▶ **Symmetry principles** brought order in the chaos:
- ▶ three quarks (**up, down, strange**)
- ▶ Murray Gell-Mann
Nobel Prize in Physics (1969)




The Eightfold Way

- ▶ **symmetry** two quantum numbers: **Isospin** and **Strangeness**
- ▶ **Isospin and Strangeness conserved** in strong interactions





The Eightfold Way

- **Mesons:** “add” a quark and an anti-quark (ex: $|\pi^- \rangle = |d\bar{u}\rangle$)



The Eightfold Way

- ▶ **Mesons:** “add” a quark and an anti-quark (ex: $|\pi^- \rangle = |d\bar{u}\rangle$)
- ▶ **Baryons:** “add” three quarks (ex: $|p\rangle = |uud\rangle$)



Color

- ▶ Trouble: get only all observed hadrons if one puts three quarks in the same state!
- ▶ BUT: quarks must have spin 1/2
- ▶ they must be fermions
(who don't like to be in the same room of the fermion motel!)
- ▶ but the model works:
predicted $|\Omega^- \rangle = |sss \rangle$ was found

Color


- ▶ Trouble: get only all observed hadrons if one puts **three quarks in the same state!**
- ▶ BUT: quarks must have **spin 1/2**
- ▶ they must be **fermions**
(**who don't like to be in the same room of the fermion motel!**)
- ▶ but the model works:
predicted $|\Omega^-\rangle = |sss\rangle$ was found
- ▶ Solution: Each quark comes in three "**colors**"
- ▶ All quarks of the same kind are the same except they can differ in the color quantum number \Rightarrow **Symmetry!**

Quantum Chromo Dynamics


- ▶ More trouble: Nobody has seen **free quarks** yet!

I want free quarks!

⇒ **break up a meson**




cannot break the meson, but I
produce more hadrons!



- ▶ quarks **confined** in hadrons
- ▶ 1973: Gross and Wilczek, Politzer
- ▶ build theory based on **color symmetry!**
- ▶ force **becomes stronger** for longer distances
- ▶ reason: force carriers themselves have **color**

Quantum Chromo Dynamics

- ▶ from color **symmetry** of quarks (color charge **conserved**)
- ▶ force carriers: **gluons** (spin 1)
- ▶ matter particles: **quarks** (spin 1/2)
- ▶ theory called **Quantum Chromo Dynamics (QCD)**
(Greek: chromos=color)
- ▶ force becomes weaker at small distances/high energy




Nobel prize in physics 2004:



Gross, Wilczek, Politzer


Reminder: Mass problem

- ▶ atomic nucleus is composed of **nucleons=protons and neutrons**
- ▶ nucleons are composed of **up** and **down** quarks



- ▶ **up** quark: charge $+2/3$, mass $m_u = 3 \text{ MeV}/c^2$
down quark: charge $-1/3$, mass $m_d = 6 \text{ MeV}/c^2$
electron: charge -1 , mass $m_e = 0.5 \text{ MeV}/c^2$
- ▶ **BUT:** nucleon mass $m_p = m_n = 940 \text{ MeV}/c^2$

Constituent Quarks




- ▶ Quarks inside hadron
- ▶ cloud of gluons around
- ▶ effectively: extended object
- ▶ constituent quark
- ▶ $M_{\text{con}} = 350 \text{ MeV}/c^2$
- ▶ $m_d = 6 \text{ MeV}/c$,
 $m_u = 3 \text{ MeV}/c$

Where does the Constituent-Quark Mass come from?

Mass generation


- ▶ Strong force at low energies very strong \Rightarrow forms $q\bar{q}$ pairs
- ▶ pairs are **bosons!** \Rightarrow all like to stay in **vacuum state** (at lowest possible energy)
- ▶ about 5 pairs per fm^{-3} (1 fm = 10^{-15} m!)




- ▶ Quarks become massive, because of very dense vacuum!
- ▶ How can we check this?
- ▶ Can we evaporate the vacuum?

How to evaporate the vacuum?

- ▶ lots of **quarks** and **gluons** close together
- ▶ **dense and hot environment** \Rightarrow strong force becomes weaker!
- ▶ **QCD** at high temperatures and densities
- ▶ $\bar{q}q$ condensate dissolves (**phase transition!**)




Use our favorite tool: Heavy-Ion Colliders!




- ▶ RHIC: Accelerate **gold nuclei** to 200 GeV per nucleon
- ▶ collide them head-on!
- ▶ Hope to create the **Quark-Gluon Plasma!**

The “Little Bang” in the Lab




- ▶ What are probes from hot and dense stage?

The “Little Bang” in the Lab



- ▶ What are probes from **hot and dense stage**?
- ▶ Answer: decays of $\rho(770)$ meson to electrons and positrons!



The “Little Bang” in the Lab



- ▶ Challenge: Find the rare events!
- ▶ See Prof. Mioduszewski's talk from last week!

The ρ meson in the vacuum

- ▶ mass of the ρ mesons: $m_\rho = 770 \text{ MeV}/c^2$
- ▶ $m_\rho \approx 2M_{\text{constituent quarks}}$
- ▶ its lifetime is about $1.3 \text{ fm}/c = 3.3 \cdot 10^{-24} \text{ sec}$
- ▶ **It decays inside the hot and dense matter!**




The ρ meson in the fireball

- ▶ how to measure the ρ mass inside the fireball?
 - ▶ could look at the decay pions
 - ▶ energy-momentum conservation $\Leftrightarrow \rho$ mass ($E = mc^2!$)
 - ▶ but pions interact strongly with the “junk” around them \Rightarrow
Signal gets destroyed!

The ρ meson in the fireball



- ▶ how to measure the ρ mass inside the fireball?
 - ▶ could look at the decay pions
 - ▶ energy-momentum conservation $\Leftrightarrow \rho$ mass ($E = mc^2!$)
 - ▶ but pions interact strongly with the “junk” around them \Rightarrow
Signal gets destroyed!
- ▶ solution: rarely the ρ 's decay into an e^+e^- or $\mu^+\mu^-$ pair



- ▶ e^\pm and μ^\pm are leptons
- ▶ they do not interact strongly
- ▶ signal undistorted
- ▶ get the mass of the ρ
inside the fireball

What do the Theoreticians predict?

- ▶ some theoreticians predicted “dropping ρ mass”
- ▶ quark condensate melts, not much else happens to the ρ
- ▶ others simulated interactions of the ρ in the hot gas
- ▶ ρ becomes a broad mass distribution (“melting ρ ”)




What is right?

- ▶ Only experiment can answer!
- ▶ most recent data from NA60 Experiment at CERN
- ▶ look for $\mu^+\mu^-$ pairs (“dileptons”)

What is right?


- ▶ Only experiment can answer!
- ▶ most recent data from NA60 Experiment at CERN
- ▶ look for $\mu^+\mu^-$ pairs (“dileptons”)



Curves from recent calculation
[Ralf Rapp, HvH 2006]
“Melting ρ ” favored compared
to dropping ρ mass

What is right?

- ▶ Only experiment can answer!
- ▶ most recent data from NA60 Experiment at CERN
- ▶ look for $\mu^+\mu^-$ pairs (“dileptons”)



Curves from recent calculation
[Ralf Rapp, HvH 2006]
“Melting ρ ” favored compared
to dropping ρ mass

- ▶ need more detailed theoretical studies
- ▶ going on also at Texas A&M!
- ▶ We begin to understand the origin of mass!

Summary

- ▶ Natural Laws \leftrightarrow Symmetries \leftrightarrow Conservation Laws
- ▶ Atom \rightarrow Nucleus \rightarrow Nucleons \rightarrow quarks (elementary!)
- ▶ Quarks always confined to Hadrons (baryons and mesons)
 - ▶ strong force described by QCD (based on color symmetry!)
 - ▶ Confinement only partially understood from QCD

Summary

- ▶ Natural Laws \leftrightarrow Symmetries \leftrightarrow Conservation Laws
- ▶ Atom \rightarrow Nucleus \rightarrow Nucleons \rightarrow quarks (elementary!)
- ▶ Quarks always confined to Hadrons (baryons and mesons)
 - ▶ strong force described by QCD (based on color symmetry!)
 - ▶ Confinement only partially understood from QCD
- ▶ Quarks acquire a large mass within hadrons
 - ▶ vacuum is a “dense liquid” of $\bar{q}q$ and gluon condensates
 - ▶ spontaneous symmetry breaking
 - ▶ more than 98% of the visible mass in the universe

Summary

- ▶ Natural Laws \leftrightarrow **Symmetries** \leftrightarrow **Conservation Laws**
- ▶ Atom \rightarrow Nucleus \rightarrow Nucleons \rightarrow **quarks** (elementary!)
- ▶ Quarks always **confined** to **Hadrons** (baryons and mesons)
 - ▶ strong force described by **QCD** (based on **color symmetry!**)
 - ▶ **Confinement** only partially understood from **QCD**
- ▶ **Quarks** acquire a **large mass** within hadrons
 - ▶ vacuum is a “**dense liquid**” of $\bar{q}q$ and gluon condensates
 - ▶ **spontaneous symmetry breaking**
 - ▶ more than 98% of the visible mass in the universe
- ▶ Collisions of Heavy Nuclei at High Energies
 - ▶ **Heat the vacuum** and **evaporate the condensates**
 - ▶ **deconfine** quarks and gluons
 - ▶ **dissolve mass** into energy
 - ▶ dileptons (e^+e^- or $\mu^+\mu^-$) from decays of ρ mesons
 - ▶ study the **origin of mass**

Summary

- ▶ Natural Laws \leftrightarrow **Symmetries** \leftrightarrow **Conservation Laws**
- ▶ Atom \rightarrow Nucleus \rightarrow Nucleons \rightarrow **quarks** (elementary!)
- ▶ Quarks always **confined** to **Hadrons** (baryons and mesons)
 - ▶ strong force described by **QCD** (based on **color symmetry!**)
 - ▶ **Confinement** only partially understood from **QCD**
- ▶ **Quarks** acquire a **large mass** within hadrons
 - ▶ vacuum is a “**dense liquid**” of $\bar{q}q$ and gluon condensates
 - ▶ **spontaneous symmetry breaking**
 - ▶ more than 98% of the visible mass in the universe
- ▶ Collisions of Heavy Nuclei at High Energies
 - ▶ **Heat the vacuum** and **evaporate the condensates**
 - ▶ **deconfine** quarks and gluons
 - ▶ **dissolve mass** into energy
 - ▶ dileptons (e^+e^- or $\mu^+\mu^-$) from decays of ρ mesons
 - ▶ study the **origin of mass**

Very exciting physics ahead!