


# Dileptons in Hot and/or Dense Matter and the Chiral Phase Transition

Hendrik van Hees

Justus-Liebig-Universität Gießen

February 22, 2011



**Institut für  
Theoretische Physik**



- 1 QCD and Chiral Symmetry
- 2 Chiral Symmetry and Hadron Phenomenology
- 3 Vector Mesons and electromagnetic Probes
- 4 Effective Models for Hadronic Many-Body Theory
- 5 Comparison with dilepton data@SPS and RHIC
- 6 Conclusions and Outlook

# QCD and (“accidental”) Symmetries

- Theory for strong interactions: QCD

$$\mathcal{L}_{\text{QCD}} = -\frac{1}{4}F_a^{\mu\nu}F_{\mu\nu}^a + \bar{\psi}(i\not{D} - \hat{M})\psi$$

- Particle content:
  - $\psi$ : Quarks, including flavor- and color degrees of freedom,  
 $\hat{M} = \text{diag}(m_u, m_d, m_s, \dots) =$  current quark masses
  - $A_\mu^a$ : gluons, gauge bosons of  $\text{SU}(3)_{\text{color}}$
- Symmetries
  - fundamental building block: local  $\text{SU}(3)_{\text{color}}$  symmetry
  - in light-quark sector: approximate chiral symmetry
  - dilation symmetry (scale invariance for  $M \rightarrow 0$ )

# "Fate" of Symmetries

- classical field theory: continuous symmetry  $\Rightarrow$  **conserved current**
- chiral limit:  $\hat{M} \rightarrow 0 \Rightarrow$ , scalar and pseudoscalar  $U(1)$  symmetries
  - $\psi \rightarrow \exp[-i(\alpha_s + \gamma_5 \alpha_p)]\psi$
  - scalar and pseudoscalar currents:  
 $\vec{j}_\mu^{(0)} = \bar{\psi}\gamma^\mu\psi, \quad \vec{j}_{A\mu}^{(0)} = \bar{\psi}\gamma_5\gamma^\mu\psi$
  - $U(1)_A$  does **not** survive quantization (**Anomaly**)
  - $\partial^\mu \vec{j}_{A\mu}^{(0)} = \frac{3}{8}\alpha_s \epsilon^{\mu\nu\rho\sigma} A_{\mu\nu}^a A_{\rho\sigma}^a$
  - Not a "bug" but a feature:
    - $\eta'$  **not** a (pseudo-)Goldstone boson
    - correct rate for  $\pi^0 \rightarrow 2\gamma$

Remark: Anomalies potential trouble in standard model of strong and "electroweak" interactions  $\leftrightarrow$  "cured" by particle content, because anomaly contributions from quarks and leptons cancel exactly!

# "Fate" of symmetries

- in classical field theory: each continuous symmetry defines **conserved current** (Noether's theorem)
- chiral limit:  $\hat{M} \rightarrow 0 \Rightarrow$ , vector-axial-vector symmetries
  - $\psi \rightarrow \exp[-i(\vec{\alpha}_V + \gamma_5 \vec{\alpha}_A) \vec{T}] \psi$   
 $\vec{T}$ : generators of  $SU(2)_{\text{flavor}}$  (or  $SU(3)_{\text{flavor}}$ )
  - **conserved vector and axial-vector currents:**

$$\vec{j}_V^\mu = \bar{\psi} \vec{T} \gamma^\mu \psi, \quad \vec{j}_A^\mu = \bar{\psi} \vec{T} \gamma_5 \gamma^\mu \psi$$


- in vacuum, chiral symmetry **spontaneously broken** by **quark condensate**:  
 $\langle 0 | \bar{\psi} \psi | 0 \rangle \neq 0$
- (approximate) Goldstone bosons:  $\pi, K, \eta$  (pseudoscalar octet)
- "real world": chiral symmetry slightly **explicitly broken** by quark masses  
 $\hat{M} \neq 0: \quad SU_L(2) \times SU_R(2) \Rightarrow SU_V(2)$
- **isospin symmetry** slightly broken by light-quark-mass differences

# "Fate" of Symmetries

- classical field theory: continuous symmetry  $\Rightarrow$  conserved current
- $\hat{M} \rightarrow 0 \Rightarrow$  dilatation (or scale) symmetry
  - $x \rightarrow \lambda x, \quad \psi \rightarrow \lambda^{-3/2}\psi, \quad A_\mu^a \rightarrow \lambda^{-1} A_\mu^a$
  - dilatation current:  
$$j_D^\mu = x_\nu \Theta^{\mu\nu}$$
  - Scale invariance does **not** survive quantization ("Trace" Anomaly)  
$$\partial_\mu j_D^\mu = \Theta_\mu{}^\mu = -\frac{\beta(\alpha_s)}{4\alpha_s} A_{\mu\nu}^a A^{a\mu\nu}$$
  - $\beta(\alpha_s)$ : Gell-Mann Low function, rules the running of the coupling with renormalization **scale**
  - Not a "bug" but a feature: hadrons get most of their mass from it!


# Phenomenology from Chiral Symmetry

- Use (approximate) **chiral symmetry** to build effective models
- **Ward identities**
  - PCAC:  $\langle 0 | \partial^\mu j_{A\mu}^k | \pi^j(\vec{k}) \rangle = iF_\pi m_\pi^2 \delta^{kj}$
  - $m_\pi^2 F_\pi^2 = -(m_u + m_d) \langle 0 | \bar{u}u | 0 \rangle$  (GOR relation)
- Spontaneous breaking causes splitting of chiral partners:



# Finite Temperature/Density: Idealized theory picture

- partition sum:  $Z(V, T, \mu_q, \Phi) = \text{Tr}\{\exp[-(\mathbf{H}[\Phi] - \mu_q \mathbf{N})/T]\}$



[ . . ., K. Chou, Z. Su, B. Hao, L. Yu 85, N.P. Landsmann, C.G. van Weert 87, E.A. Calzetta, B. Hu 08, . . . ]



- Asymptotic freedom  $\Rightarrow$  **quark condensate melts** at high enough **temperatures**
- all bulk properties from **partition sum**:

$$Z(V, T, \mu_q) = \text{Tr}\{\exp[-(\mathbf{H} - \mu_q \mathbf{N})/T]\}$$

- Free energy:  $\Omega = -\frac{T}{V} \ln Z = -P$
- **Quark condensate**:  $\langle \bar{\psi}_q \psi_q \rangle_{T, \mu_q} = \frac{V}{T} \frac{\partial P}{\partial m_q}$
- Lattice QCD indicates: Chiral symmetry restoration  $\leftrightarrow$  deconfinement phase transition (same  $T_c$ )

# Why Electromagnetic Probes?

- $\gamma, \ell^\pm$ : only e. m. interactions
- reflect whole “history” of collision
- chance to see chiral symm. rest. directly?





Fig. by A. Drees [R. Rapp, J. Wambach, *Adv. Nucl. Phys.* **25**, 1 (2000)]

# Vector Mesons and electromagnetic Probes

- **photon** and **dilepton** thermal emission rates given by **same electromagnetic-current-correlation function** ( $J_\mu = \sum_f Q_f \bar{\psi}_f \gamma_\mu \psi_f$ )

[L. McLerran, T. Toimela 85, H. A. Weldon 90, C. Gale, J.I. Kapusta 91]

$$\Pi_{\mu\nu}^<(q) = \int d^4x \exp(iq \cdot x) \langle J_\mu(0) J_\nu(x) \rangle_T = -2f_B(q_0) \text{Im} \Pi_{\mu\nu}^{(\text{ret})}(q)$$

$$q_0 \frac{dN_\gamma}{d^4x d^3\vec{q}} = \frac{\alpha}{2\pi^2} g^{\mu\nu} \text{Im} \Pi_{\mu\nu}^{(\text{ret})}(q) \Big|_{q_0=|\vec{q}|} f_B(q_0)$$

$$\frac{dN_{e^+e^-}}{d^4x d^4q} = -g^{\mu\nu} \frac{\alpha^2}{3q^2\pi^3} \text{Im} \Pi_{\mu\nu}^{(\text{ret})}(q) \Big|_{q^2=M_{e^+e^-}^2} f_B(q_0)$$

- to lowest order in  $\alpha$ :  $e^2 \Pi_{\mu\nu} \simeq \Sigma_{\mu\nu}^{(\gamma)}$
- **vector-meson dominance** model:

$$\Sigma_{\mu\nu}^{(\gamma)} = \text{Diagram with } G_\rho$$

- derivable from **partition sum**  $Z(V, T, \mu, \Phi)$ !

# Vector Mesons and chiral symmetry

- **vector** and **axial-vector** mesons  $\leftrightarrow$  correlators of the respective currents


$$\Pi_{V/A}^{\mu\nu}(p) := \int d^4x \exp(ipx) \left\langle J_{V/A}^{\nu}(0) J_{V/A}^{\mu}(x) \right\rangle_{\text{ret}}$$

- Ward-Takahashi Identities from chiral symmetry  $\Rightarrow$  **Weinberg-sum rules**



$$f_{\pi}^2 = - \int_0^{\infty} \frac{dp_0^2}{\pi p_0^2} [\text{Im } \Pi_V(p_0, 0) - \text{Im } \Pi_A(p_0, 0)]$$

- spectral functions of vector (e.g.  $\rho$ ) and axial vector (e.g.  $a_1$ ) directly related to **order parameters of chiral symmetry!**

# Vector Mesons and chiral symmetry



from [R. Rapp, *Pramana* **60**, 675 (2003)]



from [R. Rapp, *J. Phys. G* **31**, S217 (2005)]


- different models with chiral symmetry: equivalent only on shell (“low-energy theorems”)
- model-independent conclusions only in low-temperature/density limit (chiral perturbation theory) or from lattice-QCD calculations
- mass spectrum of vector mesons depends on realization chiral symmetry
  - hidden local symmetry [Bando, Kugo, PRL 54, 1215 (1984)]  $\Rightarrow$  “vector manifestation” of  $\chi$ S:  $m_\rho \rightarrow 0$  (“dropping mass”)
  - generalized hidden local symmetry ( $\rho$  and  $a_1$  as gauge fields): “normal realization” of  $\chi$ S:  $m_\rho \simeq \text{const}$  (“melting resonances”)
- use phenomenological hadronic models + many-body techniques to assess medium modifications of vector mesons

# QCD sum rules



$$M_n = - \int_0^\infty \frac{ds}{\pi} s^n [\text{Im } \Pi_V(s) - \text{Im } \Pi_A(s)]$$
$$M_{-2} = \frac{1}{3} f_\pi^2 \langle r_\pi^2 \rangle - F_A, \quad M_{-1} = f_\pi^2$$
$$M_0 = 0, \quad M_1 = c\alpha_s \langle (\bar{q}q)^2 \rangle$$

[Weinberg 67; Das et al 67; Kapusta, Shuryak 93]

- theory connection of **chiral symm. restoration** with dileptons in HICs
  - $\Pi_V, \Pi_A$  from **chiral hadronic model** at finite  $T, \mu_B$
  - compare  $M_n(T, \mu_B)$  to **lQCD chiral order parameters at finite  $T$**
  - compare  $\Pi_V$  from **hadronic model** to dileptons from HICs
- QCD sum rules
  - relate **current correlators** to **condensates**
  - VMD  $\Leftrightarrow$  **vector-meson spectral functions**
  - **dropping-mass** and **broadening scenarios** possible



- Phenomenological hadronic models [Chanfray et al, Herrmann et al, Rapp, Wambach et al, . . .] for vector mesons
- important ingredients:  $\pi\pi$  interactions  
baryonic excitations






- Baryon (resonances)** important, even at RHIC with low **net** baryon density  $n_B - n_{\bar{B}}$
- reason:  $n_B + n_{\bar{B}}$  relevant (CP invariance of strong interactions)



# The $\rho$ -meson (vacuum)





- most important for  $\rho$ -meson: pions




# The $\rho$ -meson (cold matter)

[W. Peters, M. Post, H. Lenske, S. Leupold, U. Mosel, NPA 632, 109 (1998)]

- include all four-star resonances up to  $M = 1.9$  GeV
- particularly important: **s-wave coupling**  $N(1520)$ ,  $\Delta(1620)$ ,  $\Delta(1650)$ ,  $\Delta(1700)$



# Dilepton rates: Hadron gas $\leftrightarrow$ QGP



- at  $T \simeq T_c$ : HG  $\simeq$  QGP
- QGP rate
  - HTL improved  $\bar{q} + q \rightarrow l^+ + l^-$
  - in good agreement with IQCD results
  - similar results also for  $\gamma$  rates
- “quark-hadron duality”?

[R. Rapp, J. Wambach, Eur. Phys. J. A 6, 415 (1999)]

# Sources of dilepton emission in heavy-ion collisions

- ① “core”  $\Leftrightarrow$  emission from thermal source [McLerran, Toimela 1985]

$$\frac{1}{q_T} \frac{dN^{(\text{thermal})}}{dM dq_T} = \int d^4x \int dy \int M d\varphi \frac{dN^{(\text{thermal})}}{d^4x d^4q} \text{Acc}(M, q_T, y)$$

- ② initial hard processes: Drell Yan
- ③ “corona”  $\Leftrightarrow$  emission from “primordial” mesons (jet-quenching)
- ④ after thermal freeze-out  $\Leftrightarrow$  emission from “freeze-out” mesons



[Cooper, Frye 1975]

$$N^{(\text{fo})} = \int \frac{d^3q}{q_0} \int q_\mu d\sigma^\mu f_B(u_\mu q^\mu / T) \frac{\Gamma_{\text{meson} \rightarrow \ell^+ \ell^-}}{\Gamma_{\text{meson}}} \text{Acc}$$

- use simple homogeneous cylindrical fireball of thermalized medium


# CERES/NA45 dielectron spectra

- good agreement also for **dielectron** spectra in 158 GeV Pb-Au
- **low-mass tail** from **baryon** effects



# M spectra (in $p_T$ slices)


- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )



[HvH, R. Rapp, NPA 806, 339 (2008)]

# M spectra (in $p_T$ slices)


- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )



[HvH, R. Rapp, NPA 806, 339 (2008)]

# M spectra (in $p_T$ slices)

- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )




[HvH, R. Rapp, NPA 806, 339 (2008)]



# M spectra (in $p_T$ slices)


- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )



[HvH, R. Rapp, NPA 806, 339 (2008)]

# M spectra (in $p_T$ slices)


- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )



[HvH, R. Rapp, NPA 806, 339 (2008)]

# M spectra (in $p_T$ slices)


- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )



[HvH, R. Rapp, NPA 806, 339 (2008)]

# M spectra (in $p_T$ slices)


- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )



[HvH, R. Rapp, NPA 806, 339 (2008)]

# M spectra (in $p_T$ slices)


- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )



[HvH, R. Rapp, NPA 806, 339 (2008)]

# M spectra (in $p_T$ slices)


- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )



[HvH, R. Rapp, NPA 806, 339 (2008)]

# M spectra (in $p_T$ slices)


- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )



[HvH, R. Rapp, NPA 806, 339 (2008)]

# M spectra (in $p_T$ slices)

- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )




[HvH, R. Rapp, NPA 806, 339 (2008)]



# M spectra (in $p_T$ slices)


- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )



[HvH, R. Rapp, NPA 806, 339 (2008)]

# M spectra (in $p_T$ slices)


- norm corrected by  $\sim 3\%$  due to centrality correction  
(min-bias data:  $\langle N_{\text{ch}} \rangle = 120$ , calculation  $N_{\text{ch}} = 140$ )



[HvH, R. Rapp, NPA 806, 339 (2008)]


# Importance of baryon effects

- Baryonic interactions important!
- in-medium broadening
- low-mass tail!




# Dileptons@RHIC: New Puzzle?

- huge enhancement in the LMR unexplained yet!



QGP+HMBT: R. Rapp, HvH

[A. Adare et al (PHENIX), arXiv:0912.0244 [nucl-ex]]



HSD: Linnyk, Bratkovskaya, Cassing

[E. L. Bratkovskaya, O. Linnyk, W. Cassing, J. Phys. Conf.Ser. **230**, 012032 (2010)]

# Ulrich's Legacy in "Dilepton Physics"

- **early 1990ies:** Low-energy  $NA$  and  $AA$ 
  - OBE models, form factors, gauge invariance,...
  - BUU approach for  $N$ ,  $\Delta$ ,  $N^*(1535)$ ,  $N^*(1440)$ ,  $\pi$ ,  $\eta$
  - **dileptons** including **medium modifications of  $\rho$ 's**
  - **Collaborators:** M. Schäfer, H. C. Dönges, A. Engel, G. Wolf, W. Cassing, W. Ehehalt, E. Bratkovskaya, F. de Jong...
- **late 1990ies:** in-medium spectral properties of  $\rho$ 's, relativistic HICs, QGP
  - further studies on OBE models, bremsstrahlung, formfactors,... (see T. Galatyuk's talk)
  - $\rho$  mesons in nuclear matter, importance of baryon resonances ( $p$ - and  **$s$ -waves**)
  - QCD sum rules
  - dilepton emission from **QGP** (survival of  $\rho$ -like excitations!)
  - **Collaborators:** R. Shyam, W. Peters, M. Post, H. Lenske, S. Leupold, M. Thoma,...

# Conclusions and Outlook

- dilepton spectra  $\Leftrightarrow$  in-medium em. current correlator
  - insight into fundamental (symmetry) properties of QCD
  - properties of hot/dense strongly interacting matter  $\Leftrightarrow$  QCD-phase diagram
  - chiral symmetry (restoration)
  - origin of hadron mass?!?
- model for dilepton sources
  - radiation from thermal sources: QGP,  $\rho$ ,  $\omega$ ,  $\phi$
  - $\rho$ -decay after thermal freeze-out
  - decays of non-thermalized primordial  $\rho$ 's
  - Drell-Yan annihilation, correlated  $D\bar{D}$  decays
- invariant-mass spectra and medium effects
  - excess yield dominated by radiation from thermal sources
    - baryons essential for in-medium properties of vector mesons
    - melting  $\rho$  with little mass shift robust signal! (independent of  $T_c$ )
    - "parton-hadron" duality of rates
    - $\Leftrightarrow$  compatible with chiral-symmetry restoration!
  - dimuons in In-In (NA60), Pb-Au (CERES/NA45),  $\gamma$  in Pb-Pb (WA98)

- fireball/freeze-out dynamics  $\Leftrightarrow m_T$  spectra and effective slopes
  - “non-thermal sources” important for  $q_T \gtrsim 1$  GeV
  - lower  $T_c \Rightarrow$  higher hadronic temperatures  $\Rightarrow$  harder  $q_T$  spectra
  - to describe measured effective slopes  $a_{\perp} = 0.085c^2/\text{fm} \rightarrow 0.1c^2/\text{fm}$
  - off-equilibrium effects (viscous hydro)?
- Further developments
  - understand recent PHENIX results (large dilepton excess in LMR)
  - vector- should be complemented with axial-vector-spectral functions ( $a_1$  as chiral partner of  $\rho$ )
  - constrained with IQCD via in-medium Weinberg chiral sum rules
  - direct connection to chiral phase transition!
- recent review: [R. Rapp, J. Wambach, HvH., Landolt-Brnstein, Volume I/23, 4-1 (2010)]