

## Exercise Sheet 12

### Problem 1: Spontaneously breaking discrete and continuous symmetries

Consider the following Lagrangian

$$\begin{aligned}\mathcal{L} &= \frac{1}{2}(\partial_\mu\phi)^2 - V(\phi) \\ &= \frac{1}{2}(\partial_\mu\phi)^2 - \frac{1}{2}\mu^2\phi^2 - \frac{1}{4}\lambda\phi^4\end{aligned}\quad (1)$$

that describes a real scalar field  $\phi$  with a specific potential term. It is straightforward to show that  $\mathcal{L}$  is invariant under  $\phi \rightarrow -\phi$ . Regarding the parameters  $(\lambda, \mu^2)$ ,  $\lambda$  should be positive to ensure the existence of a ground state but, in principle,  $\mu^2$  could be positive or negative. The minimum of  $V(\phi)$ ,  $\phi_0$ , turns out to be:

$$\phi_0 = \begin{cases} 0 & \text{for } \mu^2 > 0 \\ \pm\sqrt{\frac{-\mu^2}{\lambda}} \equiv \nu & \text{for } \mu^2 < 0 \end{cases}\quad (2)$$




Figure 1:  $V(\phi)$  as given by Eq. 1 in two different cases:  $\mu^2 > 0$  (left) and  $\mu^2 < 0$  (right).

The potential,  $V(\phi)$ , in both scenarios is shown in Fig. 1. For  $\mu^2 > 0$  the quantum field must have a null vacuum expectation value (VEV). This is not the case for  $\phi$  when  $\mu^2 < 0$

$$\langle 0|\phi|0\rangle = \nu \quad (3)$$

i.e. the vacuum is degenerate. Thus, a field  $\eta$  that is centered at the vacuum should be introduced

$$\eta = \phi - \nu \quad (4)$$

such that this shifted field satisfies

$$\langle 0|\eta|0\rangle = 0. \quad (5)$$

Then, at the quantum level, the same system is described by  $\eta(x)$  with Lagrangian

$$\mathcal{L} = \frac{1}{2}(\partial_\mu\eta)(\partial^\mu\eta) - \lambda\nu^2\eta^2 - \lambda\nu\eta^3 - \frac{\lambda}{4}\eta^4 \quad (6)$$

that describes a massive scalar field ( $m_\eta = \sqrt{2\lambda\nu}$ ) but is no longer invariant under  $\eta \rightarrow -\eta$ .  $\mathcal{L}(\phi)$  had the symmetry but the parameters ( $\lambda, \mu^2$ ) can be such that the ground state of the Hamiltonian is not symmetric: the symmetry has been spontaneously broken.

The same procedure can be applied to more complex scenarios.

Consider a complex scalar field  $\phi(x)$  with Lagrangian

$$\mathcal{L} = (\partial_\mu \phi^*)(\partial^\mu \phi) - V(|\phi|) = (\partial_\mu \phi^*)(\partial^\mu \phi) - \mu^2 |\phi|^2 - \lambda |\phi|^4 \quad (7)$$

Show that it is invariant under a global U(1) symmetry, i.e.,  $\phi \rightarrow \exp(-i\alpha)\phi$ ,  $\alpha = \text{const} \in \mathbb{R}$ . Write down the Lagrangian in terms of  $(\phi_1, \phi_2)$  where

$$\phi = \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2) \quad (8)$$

Repeat the previous procedure in the case of the discrete symmetry to compute  $\phi_0$ . Write down the Lagrangian in terms of the shifted fields:

$$\begin{aligned} \eta &= \phi_1 - \nu \\ \xi &= \phi_2 \end{aligned} \quad (9)$$

How many massive and massless scalar particles does it contain? What does this fact have to do with the broken symmetry?

**Hint:** If we repeat the same calculation but with an SU(2) triplet instead of a scalar, the Lagrangian in terms of the shifted fields would no longer be invariant under SU(2) but under U(1) and it would contain 2 massless scalar fields.

---

## Problem 2: Spontaneously breaking gauge invariance

Consider a U(1) gauge-invariant Lagrangian for a complex scalar field  $\phi(x)$

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + (D_\mu \phi)^*(D^\mu \phi) - \underbrace{\mu^2 \phi^* \phi - \lambda(\phi^* \phi)^2}_{-V(|\phi|)} \quad (10)$$

with the gauge-covariant derivative

$$D_\mu = \partial_\mu + iqA_\mu(\underline{x}) \quad (11)$$

and the field-strength tensor

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \quad (12)$$

that is invariant under

$$\begin{aligned} \phi'(\underline{x}) &= \exp[-iq\Theta(\underline{x})]\phi(\underline{x}) \\ A'_\mu(\underline{x}) &= A_\mu(x) + \partial_\mu \Theta(x) \end{aligned} \quad (13)$$

Compute the minimum  $\phi_0 = \nu/\sqrt{2} > 0 \in \mathbb{R}$  of the potential of  $V(|\phi|)$  for  $\mu^2 < 0$ . In this case instead of writing  $\phi$  in terms of the shifted fields  $(\eta, \xi)$  as

$$\phi(x) \equiv \frac{1}{\sqrt{2}}[\nu + \eta(x) + i\xi(x)], \quad \langle 0|\eta|0\rangle = \langle 0|\xi|0\rangle = 0 \quad (14)$$

it is more transparent to write

$$\phi(x) \equiv \exp(-iq\xi(x)/\nu) \frac{1}{\sqrt{2}}[\nu + \eta(x)], \quad \langle 0|\eta|0\rangle = \langle 0|\xi|0\rangle = 0 \quad (15)$$

and exploit local gauge invariance so that

$$\phi'(x) = \exp(iq\xi(x)/\nu)\phi(x) = \frac{1}{\sqrt{2}}[\nu + \eta(x)] \quad (16)$$

that is equivalent to choose the gauge field  $\Theta(x) = -\xi(x)/\nu$ . This choice is called unitary gauge. Rewrite  $\mathcal{L}$  in terms of  $\eta(x)$  and analyse the different terms that it contains. Is the gauge boson massless?

Discuss, how the various “field-degrees of freedom” are redistributed when comparing the cases with and without broken *local* gauge invariance, and discuss, why there are no massless Goldstone modes in this case in contradistinction to the case of a spontaneously broken *global* symmetry, as considered in Problem 1.