

## Exercise Sheet #7

Fabian Schubert <fschubert@itp.uni-frankfurt.de>  
Oren Neumann <neumann@itp.uni-frankfurt.de>

**Problem 1** (*Hausdorff Dimension*) 8 Pts

Calculate the Hausdorff dimension of a straight line and of the Cantor set.

The Cantor ternary set is created by repeatedly deleting the open middle thirds of a set of line segments. First, one removes the open middle third  $(1/3, 2/3)$  from the interval  $[0, 1]$ , leaving two line segments:  $[0, 1/3]$  and  $[2/3, 1]$ . Then, the open middle third of each of these remaining segments is also removed, leaving four line segments:  $[0, 1/9]$ ,  $[2/9, 1/3]$ ,  $[2/3, 7/9]$  and  $[8/9, 1]$ . Then the remaining four segments get their open middle third removed and the process is repeated infinitely.

**Problem 2** (*Lorenz system*) 12 Pts

(a) Analyse the fixpoints of the Lorenz system given by

$$\begin{aligned}\dot{x} &= s(y - x) \\ \dot{y} &= x(r - z) - y \\ \dot{z} &= xy - bz\end{aligned}$$

and their stability as a function of the Rayleigh number  $r > 0$  with parameters  $s = 10$  and  $b = 8/3$  fixed. Make a sketch of the  $x(r)$  bifurcation diagram. (*Hint*: You may calculate the the eigenvalues numerically.)

- (b) Show analytically that all fixed points are unstable for  $r > r_H = s(s + b + 3)/(s - b - 1)$ . (*Hint*: You do not necessarily need a full solution for the eigenvalue spectra, since you are only looking for this particular transition point.)
- (c) What kind of bifurcation does the system undergo at  $r = r_H$  and what kind of dynamics are present in the system for  $r > r_H$ ?