

Heavy Quarks in the Quark-Gluon Plasma

Hendrik van Hees

Justus-Liebig Universität Gießen

December 2, 2008

with

M. Mannarelli, V. Greco, L. Ravagli and R. Rapp



**Institut für
Theoretische Physik**

JUSTUS-LIEBIG-
 UNIVERSITÄT
GIESSEN

Outline

1 Heavy-quark interactions in the sQGP

- Heavy-quark diffusion: The Fokker-Planck Equation
- Elastic pQCD heavy-quark scattering
- Non-perturbative interactions: effective resonance model

2 Non-photonic electrons at RHIC


3 Microscopic model for non-pert. HQ interactions

- Static heavy-quark potentials from lattice QCD
- T-matrix approach


4 Resonance-Recombination Model

5 Summary and Outlook


Heavy Quarks in Heavy-Ion collisions



hard production of HQs
described by PDF's + pQCD (PYTHIA)




HQ rescattering in QGP: Langevin simulation
drag and diffusion coefficients from
microscopic model for HQ interactions in the sQGP



Hadronization to D, B mesons via
quark coalescence + fragmentation

V. Greco, C. M. Ko, R. Rapp, PLB **595**, 202 (2004)



semileptonic decay \Rightarrow
“non-photonic” electron observables

Heavy-Quark diffusion

- Fokker-Planck Equation

$$\frac{\partial f(t, \vec{p})}{\partial t} = \frac{\partial}{\partial p_i} \left[p_i \textcolor{red}{A}(t, p) + \frac{\partial}{\partial p_j} \textcolor{green}{B}_{ij}(t, \vec{p}) \right] f(t, \vec{p})$$

- drag (friction) and diffusion coefficients

$$p_i A(t, \vec{p}) = \langle p_i - p'_i \rangle$$

$$B_{ij}(t, \vec{p}) = \frac{1}{2} \langle (p_i - p'_i)(p_j - p'_j) \rangle$$

$$= B_0(t, p) \left(\delta_{ij} - \frac{p_i p_j}{p^2} \right) + B_1(t, p) \frac{p_i p_j}{p^2}$$

- transport coefficients defined via \mathcal{M}

$$\begin{aligned} \langle X(\vec{p}') \rangle &= \frac{1}{\gamma_c} \frac{1}{2E_p} \int \frac{d^3 \vec{q}}{(2\pi)^3 2E_q} \int \frac{d^3 \vec{q}'}{(2\pi)^3 2E_{q'}} \int \frac{d^3 \vec{p}'}{(2\pi)^3 2E_{p'}} \\ &\quad \sum |\mathcal{M}|^2 (2\pi)^4 \delta^{(4)}(p + q - p' - q') \hat{f}(\vec{q}) X(\vec{p}') \end{aligned}$$

- correct equil. lim. \Rightarrow Einstein relation: $B_1(t, p) = T(t) E_p A(t, p)$

Meaning of Fokker-Planck coefficients

- non-relativistic equation with constant $A = \gamma$ and $B_0 = D_1 = D$

$$\frac{\partial f}{\partial t} = \gamma \frac{\partial}{\partial \vec{p}} (\vec{p} f) + D \frac{\partial^2 f}{\partial \vec{p}^2}$$

- Green's function:

$$G(t, \vec{p}; \vec{p}_0) = \left\{ \frac{\gamma}{2\pi D [1 - \exp(-2\gamma t)]} \right\}^{3/2} \times \exp \left\{ -\frac{\gamma}{2D} \frac{[\vec{p} - \vec{p}_0 \exp(-\gamma t)]^2}{1 - \exp(-2\gamma t)} \right\}$$

- Gaussian with

$$\langle \vec{p}(t) \rangle = \vec{p}_0 \exp(-\gamma t),$$

$$\langle \vec{p}^2(t) \rangle - \langle \vec{p}(t) \rangle^2 = \frac{3D}{\gamma} [1 - \exp(-2\gamma t)] \underset{t \rightarrow 0}{\cong} 6Dt$$

- γ : friction (drag) coefficient; D : diffusion coefficient
- equilibrium limit for $t \rightarrow \infty$: $D = mT\gamma$
(Einstein's dissipation-fluctuation relation)

Relativistic Langevin process

- Fokker-Planck equation equivalent to **stochastic differential equation**
- **Langevin process:** friction force + Gaussian random force
- in the (local) rest frame of the heat bath


$$d\vec{x} = \frac{\vec{p}}{E_p} dt,$$

$$d\vec{p} = -A \vec{p} dt + \sqrt{2dt} [\sqrt{B_0} P_{\perp} + \sqrt{B_1} P_{\parallel}] \vec{w}$$

- \vec{w} : normal-distributed random variable
- dependent on **realization of stochastic process**
- to guarantee correct equilibrium limit: Use **Hänggi-Klimontovich calculus**, i.e., use $B_{0/1}(t, \vec{p} + d\vec{p})$
- for constant coefficients: Einstein dissipation-fluctuation relation
 $B_0 = B_1 = E_p T A$.
- to implement flow of the medium
 - use **Lorentz** boost to change into local “heat-bath frame”
 - use **update rule** in heat-bath frame
 - boost back into “lab frame”

Elastic pQCD processes

- Lowest-order matrix elements [Combridge 79]



- **Debye-screening mass** for *t*-channel gluon exch. $\mu_g = gT$, $\alpha_s = 0.4$
- not sufficient to understand RHIC data on “non-photonic” electrons

Non-perturbative interactions: effective resonance model

- General idea: Survival of D - and B -meson like **resonances** above T_c
- Chiral symmetry** $SU_V(2) \otimes SU_A(2)$ in light-quark sector of **QCD**

$$\mathcal{L}_D^{(0)} = \sum_{i=1}^2 [(\partial_\mu \Phi_i^\dagger)(\partial^\mu \Phi_i) - m_D^2 \Phi_i^\dagger \Phi_i] + \text{massive (pseudo-)vectors } D^*$$

- Φ_i : two doublets: **pseudo-scalar** $\sim \begin{pmatrix} \overline{D^0} \\ D^- \end{pmatrix}$ and **scalar**
- Φ_i^* : two doublets: **vector** $\sim \begin{pmatrix} \overline{D^{0*}} \\ D^{-*} \end{pmatrix}$ and **pseudo-vector**

$$\mathcal{L}_{qc}^{(0)} = \bar{q} i \not{\partial} q + \bar{c} (i \not{\partial} - m_c) c$$

- q : light-quark doublet $\sim \begin{pmatrix} u \\ d \end{pmatrix}$
- c : singlet

Interactions


- Interactions determined by **chiral** symmetry
- For transversality of vector mesons:
heavy-quark effective theory vertices

$$\begin{aligned}\mathcal{L}_{\text{int}} = & -G_S \left(\bar{q} \frac{1 + \not{v}}{2} \Phi_1 c_v + \bar{q} \frac{1 + \not{v}}{2} i \gamma^5 \Phi_2 c_v + h.c. \right) \\ & - G_V \left(\bar{q} \frac{1 + \not{v}}{2} \gamma^\mu \Phi_{1\mu}^* c_v + \bar{q} \frac{1 + \not{v}}{2} i \gamma^\mu \gamma^5 \Phi_{2\mu}^* c_v + h.c. \right)\end{aligned}$$


- v : four velocity of heavy quark
- in **HQET**: spin symmetry $\Rightarrow G_S = G_V$

Resonance Scattering

- elastic heavy-light-(anti-)quark scattering




- D - and B -meson like resonances in sQGP



- parameters

- $m_D = 2 \text{ GeV}, \Gamma_D = 0.4 \dots 0.75 \text{ GeV}$
- $m_B = 5 \text{ GeV}, \Gamma_B = 0.4 \dots 0.75 \text{ GeV}$



Cross sections



- total pQCD and resonance cross sections: comparable in size
- BUT pQCD forward peaked \leftrightarrow resonance isotropic
- resonance scattering more effective for friction and diffusion

Transport coefficients: pQCD vs. resonance scattering



- three-momentum dependence



- resonance contributions factor $\sim 2 \dots 3$ higher than pQCD!

Transport coefficients: pQCD vs. resonance scattering

- Temperature dependence



Time evolution of the fire ball

- Elliptic **fire-ball** parameterization
fitted to hydrodynamical flow pattern [Kolb '00]

$$V(t) = \pi(z_0 + v_z t)a(t)b(t), \quad a, b: \text{half-axes of ellipse},$$
$$v_{a,b} = v_\infty[1 - \exp(-\alpha t)] \mp \Delta v[1 - \exp(-\beta t)]$$



- Isentropic expansion: $S = \text{const}$ (fixed from N_{ch})
- QGP Equation of state:

$$s = \frac{S}{V(t)} = \frac{4\pi^2}{90} T^3 (16 + 10.5 n_f^*), \quad n_f^* = 2.5$$


- obtain $T(t) \Rightarrow A(t, p)$, $B_0(t, p)$ and $B_1 = TEA$
- for semicentral collisions ($b = 7 \text{ fm}$): $T_0 = 340 \text{ MeV}$,
QGP lifetime $\simeq 5 \text{ fm}/c$.
- simulate FP equation as **relativistic Langevin process**

Initial conditions


- need initial p_T -spectra of **charm** and **bottom** quarks
 - (modified) PYTHIA to describe exp. D meson spectra, assuming **δ -function fragmentation**
 - exp. **non-photonic single- e^\pm** spectra: Fix bottom/charm ratio



Spectra and elliptic flow for heavy quarks



Au-Au $\sqrt{s}=200$ GeV ($b=7$ fm)



- $\mu_D = gT, \alpha_s = g^2/(4\pi) = 0.4$
- resonances $\Rightarrow c$ -quark thermalization without upscaling of cross sections
- Fireball parametrization consistent with hydro



- $\mu_D = 1.5 T$ fixed

- spatial diff. coefficient:

$$D = D_s = \frac{T}{mA}$$



- $2\pi TD \simeq \frac{3}{2\alpha_s^2}$

Spectra and elliptic flow for heavy quarks



Observables: p_T -spectra (R_{AA}), v_2



- Hadronization: Coalescence with light quarks + fragmentation
 $\Leftrightarrow c\bar{c}, b\bar{b}$ conserved
- single electrons from decay of D - and B -mesons



- Without further adjustments: data quite well described
[HvH, V. Greco, R. Rapp, Phys. Rev. C **73**, 034913 (2006)]

Observables: p_T -spectra (R_{AA}), v_2


- Hadronization: Fragmentation only
- single electrons from decay of D - and B -mesons




Observables: p_T -spectra (R_{AA}), v_2

- Central Collisions
- single electrons from decay of D - and B -mesons


Coalescence+Fragmentation




Fragmentation only



Comparison to newer data



Microscopic model: Static potentials from lattice QCD



- color-singlet free energy from lattice
- use **internal energy**

$$U_1(r, T) = F_1(r, T) - T \frac{\partial F_1(r, T)}{\partial T},$$


$$V_1(r, T) = U_1(r, T) - U_1(r \rightarrow \infty, T)$$

- Casimir scaling for other color channels [Nakamura et al 05; Döring et al 07]

$$V_{\bar{3}} = \frac{1}{2} V_1, \quad V_6 = -\frac{1}{4} V_1, \quad V_8 = -\frac{1}{8} V_1$$

T-matrix



- Brueckner many-body approach for elastic $Qq, Q\bar{q}$ scattering



- reduction scheme: 4D Bethe-Salpeter → 3D Lipmann-Schwinger
- S - and P waves
- same scheme for light quarks (self consistent!)
- Relation to invariant matrix elements



$$\sum |M(s)|^2 \propto \sum_q d_a (|T_{a,l=0}(s)|^2 + 3|T_{a,l=1}(s)|^2 \cos \theta_{cm})$$

T-matrix



- **resonance formation** at lower temperatures $T \simeq T_c$
- **melting** of resonances at higher $T!$ \Rightarrow sQGP
- P wave smaller
- resonances near T_c : natural connection to quark coalescence
 [Ravagli, Rapp 07]
- model-independent assessment of elastic $Qq, Q\bar{q}$ scattering
- problems: uncertainties in extracting potential from IQCD
 in-medium potential V vs. F ?

Transport coefficients





- from non-pert. interactions reach $A_{\text{non-pert}} \simeq 1/(7 \text{ fm}/c) \simeq 4A_{\text{pQCD}}$
- A decreases with higher temperature
- higher density (over)compensated by melting of resonances!
- spatial diffusion coefficient

$$D_s = \frac{T}{mA}$$

increases with temperature

Non-photonic electrons at RHIC

- same model for bottom
- quark coalescence+fragmentation $\rightarrow D/B \rightarrow e + X$





- coalescence crucial for explanation of data
- increases both, R_{AA} and $v_2 \Leftrightarrow$ “momentum kick” from light quarks!
- “resonance formation” towards $T_c \Rightarrow$ coalescence natural [Ravagli, Rapp 07]

Properties of the sQGP

- measure for coupling strength in plasma: η/s
- relation to spatial diffusion coefficient

$$\frac{\eta}{s} \simeq \frac{1}{2} TD_s \quad (\text{AdS/CFT}), \quad \frac{\eta}{s} \simeq \frac{1}{5} TD_s \quad (\text{wQGP})$$



Resonance-Recombination Model


- successes of quark-coalescence models in HI phenomenology
 - high baryon/meson ration in heavy-ion compared to pp collisions compared
 - Constituent-quark number scaling of v_2
- $$v_{2,\text{had}}(p_T) = n_q v_{2,q}(p_T/n_q)$$
- experiment: CQNS better for KE_T than p_t
- problems with “naive” coalescence models
 - violates **conservation laws** (energy, momentum!)
 - violates **2nd theorem of thermodynamics** (entropy)
- **Resonance structures close to T_c**
 - transport process with $q\bar{q}(qq) \leftrightarrow R$

Resonance-Recombination Model

$$\frac{\partial}{\partial t} f_{\textcolor{red}{M}}(t, p) = -\frac{\Gamma}{\gamma_p} f_{\textcolor{red}{M}}(t, p) + g(p) \Rightarrow f_{\textcolor{red}{M}}^{(\text{eq})}(p) = \frac{\gamma_p}{\Gamma} g(p)$$



$$g(p) = \int \frac{d^3 p_1 d^3 p_2}{(2\pi)^6} \int d^3 x f_{\textcolor{blue}{q}}(x, p_1) f_{\bar{q}}(x, p_2) \sigma(s) v_{\text{rel}} \delta^{(3)}(p - p_1 - p_2)$$

$$\sigma(s) = g_\sigma \frac{4\pi}{k_{\text{cm}}^2} \frac{(\Gamma m)^2}{(s - m^2)^2 + (\Gamma m)^2}$$



Meson spectra

- $q\bar{q}$ input: Langevin simulation
- meson output: resonance-recombination model





Constituent-quark number scaling

- usual coalescence models: **factorization ansatz**

$$f_q(p, x, \varphi) = f_q(p, x)[1 + 2v_2^q(p_T) \cos(2\varphi)]$$

- CQNS usually not robust with more realistic parametrizations of v_2
- here: *q* input from Langevin simulation



Summary and Outlook

- Summary
 - Heavy quarks in the sQGP
 - non-perturbative interactions
 - mechanism for strong coupling: resonance formation at $T \gtrsim T_c$
 - IQCD potentials parameter free
 - res. melt at higher temperatures \Leftrightarrow consistency betw. R_{AA} and v_2 !
 - also provides “natural” mechanism for quark coalescence
 - resonance-recombination model
 - problems
 - extraction of V from lattice data
 - potential approach at finite T : F , V or combination?
- Outlook
 - include inelastic heavy-quark processes (gluon-radiation processes)
 - other heavy-quark observables like charmonium suppression/regeneration