

# $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$ from a momentum space analysis of the quark-antiquark static potential

Felix Karbstein<sup>1,2</sup>, Antje Peters<sup>3</sup>, Marc Wagner<sup>3,4</sup>

<sup>1</sup> Helmholtz-Institut Jena, Fröbelstieg 3, D-07743 Jena, Germany

<sup>2</sup> Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, Max-Wien-Platz 1, D-07743 Jena, Germany

<sup>3</sup> Goethe-Universität Frankfurt am Main, Institut für Theoretische Physik, Max-von-Laue-Straße 1, D-60438 Frankfurt am Main, Germany

<sup>4</sup> European Twisted Mass Collaboration (ETMC)

July 29, 2014

---

## Abstract

We determine  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$  by fitting perturbative expressions for the quark-antiquark static potential to lattice results for QCD with  $n_f = 2$  dynamical quark flavors. To this end we use the perturbative static potential at the presently best known accuracy, i.e. up to  $\mathcal{O}(\alpha_s^4)$ , in momentum space. The lattice potential is computed on a fine lattice with  $a \approx 0.042$  fm in position space. To allow for a comparison and matching of both results, the lattice potential is transformed into momentum space by means of a discrete Fourier transform. The value of  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$  is extracted in momentum space. All sources of statistical and systematic errors are discussed. The uncertainty in the value of  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$  is found to be smaller than that obtained in a recent position space analysis of the static potential based on the same lattice data. Our final results are  $r_0 \Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 0.692(31)$  and  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 331(21)$  MeV.

---

# 1 Introduction

In this paper we aim at determining  $\Lambda_{\overline{\text{MS}}}$  by comparing lattice and perturbative results for the quark-antiquark ( $Q\bar{Q}$ ) static potential<sup>1</sup> in momentum space. More precisely, we restrict ourselves to quantum chromodynamics (QCD) with  $n_f = 2$  dynamical quark flavors, i.e. exclusively focus on  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$ .

The  $Q\bar{Q}$  static potential  $V$  amounts to the interaction energy of the color-singlet state made up of a static quark  $Q$  and its antiquark  $\bar{Q}$  separated by a distance  $r = |\vec{r}|$  (in position space), or characterized by a momentum transfer  $p = |\vec{p}|$  (in momentum space), respectively. Accordingly, we denote the potential in position space by  $V(r)$  and in momentum space by  $V(p)$ .

While many studies aiming at the extraction of  $\Lambda_{\overline{\text{MS}}}$  from the  $Q\bar{Q}$  static potential have been performed so far, e.g. [1–7], to the best of our knowledge we are the first to transform the lattice data to momentum space and compare and match perturbative and lattice results in momentum space. As will be discussed in detail below, our present study is mainly motivated by the significantly worse convergence behavior of the perturbative potential in position space as compared to momentum space [8–11].

Higher-order perturbative calculations in QCD are most conveniently performed in momentum space. This is particularly true for the perturbative  $Q\bar{Q}$  static potential. Hence, the perturbative expression for the static potential at the highest current accuracy, which encompasses all contributions up to  $\mathcal{O}(\alpha_s^4)$ , is directly accessible in momentum space. Due to the fact that QCD is asymptotically free, perturbative calculations in QCD are viable only at large momentum transfers  $p \gg \Lambda_{\text{QCD}}$ , with  $\Lambda_{\text{QCD}}$  denoting the *QCD (momentum) scale*, which can be seen as separating the regimes of perturbative and non-perturbative physics. However, note that in standard perturbation theory loop diagrams come along with integrations of the loop four-momentum over the full momentum regime, implying that such loops naturally also receive contributions from momenta  $\lesssim \Lambda_{\text{QCD}}$  for which perturbation theory is no longer trustworthy. The leading uncontrolled contribution  $\delta V(p)$  contained in the perturbative potential in momentum space arising from this kind of diagrams is quadratic in  $\Lambda_{\text{QCD}}$  and scales as  $\delta V(p) \sim -\frac{4\pi\alpha_s}{p^2} \left(\frac{\Lambda_{\text{QCD}}}{p}\right)^2$  [11]. Taking into account that  $V(p) \sim -\frac{4\pi\alpha_s}{p^2} (1 + \mathcal{O}(\alpha_s))$  and  $p \gg \Lambda_{\text{QCD}}$ , i.e.  $\frac{\Lambda_{\text{QCD}}}{p} \ll 1$ , this only amounts to a tiny correction. For completeness, note that the strong coupling  $\alpha_s$  depends on an *a priori* arbitrarily chosen renormalization momentum scale  $\mu$ , i.e.  $\alpha_s = \alpha_s(\mu)$ . The  $\mu$  dependence is such that  $\alpha_s(\mu) \ll 1$  only for  $\mu \gg \Lambda_{\text{QCD}}$ . A particularly obvious choice of the renormalization scale for the couplings in  $V(p)$  is  $\mu = p$ , corresponding to an identification of  $\mu$  with the *typical momentum scale* of the quantity under consideration.

Conversely, within lattice QCD the  $Q\bar{Q}$  static potential is naturally computed in position space. It can be extracted straightforwardly by studying the exponential decay of the rectangular Wilson loop as a function of its temporal extension [12]. Of course, lattice simulations at a given lattice spacing  $a$  cannot resolve arbitrarily small separations. Moreover, the minimum attainable lattice spacing is limited by the available computing power. Additional complications in the computation of observables at small lattice spacings might arise from the problem of topology freezing [13–18]. The behavior of the static potential at small  $Q\bar{Q}$  separations is

---

<sup>1</sup>In agreement with the prevalent notation, particularly in the field of lattice QCD, we use the terms *static potential* and *static energy* synonymously. Note, however, that in the literature sometimes a distinction is made and these terms refer to different quantities.

intimately related to its behavior at large momenta. Consequently, after a Fourier transform to momentum space, lattice results are expected to allow for reliable insights only below a certain threshold momentum, which is  $\lesssim$  the maximum momentum  $\bar{p} = \pi/a$  that can be resolved on a lattice of spacing  $a$  (cf. Sec. 2 below).

Taking into account the above constraints, a comparison and matching of perturbative and lattice results in momentum space is limited to momenta  $p$  fulfilling  $\Lambda_{\text{QCD}} \ll p \ll \bar{p}$ .

Analogous considerations can be invoked to delimit the fitting interval in position space (cf. e.g. [5]), employing that the typical momentum scale that can be attributed to a relative distance  $r$  scales as  $p \sim 1/r$ . In position space the manifestly perturbative regime is thus characterized by  $1/r \gg \Lambda_{\text{QCD}}$ . However, an important difference is that the leading uncontrolled contribution  $\delta V(r)$  to the static potential in position space as determined by a standard Fourier transform from momentum to position space is more pronounced than in momentum space: it is linear in  $\Lambda_{\text{QCD}}$ , scales as  $\delta V(r) \sim \frac{\alpha_s}{r}(r\Lambda_{\text{QCD}}) = \alpha_s\Lambda_{\text{QCD}}$ , and arises exclusively from the low momentum part of the Fourier integral over momenta  $\lesssim \Lambda_{\text{QCD}}$ , for which perturbation theory is no longer trustworthy [11]<sup>2</sup>. For the position space potential a seemingly obvious choice of the renormalization scale is  $\mu = 1/r$ , which amounts to the *typical momentum scale* to be associated with the static quarks separated by a distance  $r$ .

However, it has been recognized long ago that in particular for the identification  $\mu = 1/r$  the convergence of the perturbative potential in position space as defined by a standard Fourier transform from momentum space is spoiled [19], while it can be significantly improved by subtracting just the uncontrolled contribution  $\delta V(r)$  linear in  $\Lambda_{\text{QCD}}$  [11]. Unfortunately this necessitates the specification of an additional *subtraction scale* and thereby increases the number of free parameters. Without any subtraction procedure a meaningful fit of perturbative expressions for the  $Q\bar{Q}$  static potential to lattice results in position space is not possible: the perturbative expressions only show a controlled convergence behavior for separations, which are much smaller than the minimum accessible separations on state-of-the-art lattice simulations (cf. e.g. [20–23]). Correspondingly, aiming at an accurate determination of  $\Lambda_{\overline{\text{MS}}}$  by comparing the results from lattice QCD simulations with perturbative calculations of the  $Q\bar{Q}$  static potential in position space, considerable efforts are needed to cope with this issue. Various strategies to retain or restore the significantly better convergence of  $V(p)$  also for  $V(r)$  have been devised in the literature [3, 11, 20–24].

An extraction of  $\Lambda_{\overline{\text{MS}}}$  directly in momentum space of course does not involve a Fourier transform of the perturbative potential, such that one may hope to circumvent most of these limitations. On the other hand, one now has to Fourier transform the lattice potential, which might eventually lead to similar problems. As we will argue and demonstrate in detail in this paper, most favorable for us the latter concerns are not substantiated.

Our paper is organized as follows. Section 2 is devoted to the  $Q\bar{Q}$  static potential on the lattice. After briefly reviewing its computation in position space, we point out how we transform it to momentum space. In Sec. 3 we summarize the present knowledge of the perturbative  $Q\bar{Q}$  static potential and detail on the role of  $\Lambda_{\overline{\text{MS}}}$ . Special emphasis is put on the convergence behavior of both the perturbative  $Q\bar{Q}$  static potential for  $n_f = 2$  in momentum space and the QCD

---

<sup>2</sup>Recall that in the perturbative regime both dimensionless quantities,  $\Lambda_{\text{QCD}}/p$  and  $r\Lambda_{\text{QCD}}$ , are small and of the same order of magnitude, i.e.  $\Lambda_{\text{QCD}}/p \sim r\Lambda_{\text{QCD}} \sim \epsilon \ll 1$ . Correspondingly,  $r\Lambda_{\text{QCD}} = \mathcal{O}(\epsilon)$ , while  $(\frac{\Lambda_{\text{QCD}}}{p})^2 = \mathcal{O}(\epsilon^2)$ .

$\beta$ -function. Section 4 constitutes the main section of our paper. Here we describe in detail our momentum space analysis to extract  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$  by fitting the perturbative expressions for the  $Q\bar{Q}$  static potential  $V(p)$  to the corresponding lattice results. The various error sources are identified and delineated and our final result for  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$  is specified. Moreover, comparisons with the result of [5], amounting to a position space extraction of  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$  based on the same lattice data, are made. Finally, we end with conclusions in Sec. 5.

## 2 The $Q\bar{Q}$ static potential in momentum space from lattice QCD

### 2.1 Gauge link configurations

We use the same  $n_f = 2$  gauge link configurations as for a recent determination of  $\Lambda_{\overline{\text{MS}}}$ , where, in contrast to this work, the lattice results and perturbative expressions for the static potential were compared and matched in position space [5]. These gauge link configurations were generated by the European Twisted Mass Collaboration (ETMC) [25–27] with the tree-level Symanzik improved gauge action [28],

$$S_G[U] = \frac{\beta}{6} \left( b_0 \sum_{x, \mu \neq \nu} \text{Tr} \left( 1 - P^{1 \times 1}(x; \mu, \nu) \right) + b_1 \sum_{x, \mu \neq \nu} \text{Tr} \left( 1 - P^{1 \times 2}(x; \mu, \nu) \right) \right) \quad (1)$$

with  $b_0 = 1 - 8b_1$  and  $b_1 = -1/12$  and the Wilson twisted mass quark action [29–32],

$$S_F[\chi, \bar{\chi}, U] = a^4 \sum_x \bar{\chi}(x) \left( D_W + i\mu_q \gamma_5 \tau_3 \right) \chi(x) \quad (2)$$

with

$$D_W = \frac{1}{2} \left( \gamma_\mu \left( \nabla_\mu + \nabla_\mu^* \right) - a \nabla_\mu^* \nabla_\mu \right) + m_0. \quad (3)$$

Here  $a$  denotes the lattice spacing,  $\nabla_\mu$  and  $\nabla_\mu^*$  are the gauge covariant forward and backward derivatives,  $m_0$  and  $\mu_q$  are the bare untwisted and twisted quark masses,  $\tau_3$  is the third Pauli matrix acting in flavor space, and  $\chi = (\chi^{(u)}, \chi^{(d)})$  represents the quark fields in the so-called twisted basis.

The twist angle  $\omega$  is given by  $\omega = \arctan(\mu_R/m_R)$ , where  $\mu_R$  and  $m_R$  denote the renormalized twisted and untwisted quark masses. For the ensembles of gauge link configurations considered in the present study (cf. Table 1)  $\omega$  has been tuned to  $\pi/2$  by adjusting  $m_0$  appropriately. This ensures automatic  $\mathcal{O}(a)$  improvement for many observables including the static potential (cf. [26] for details).

The considered gauge link configurations cover several different values of the lattice spacing, the pion mass  $m_{\text{PS}}$  and the spacetime volume  $L^3 \times T$ ; cf. Table 1, which also provides the number of gauge link configurations, used for the computation of the static potential, for each ensemble. The scale has been set via the pion mass and the pion decay constant, using chiral perturbation theory. Details on the generation of these gauge field configurations as well as on the computation and the analysis of standard quantities (e.g. lattice spacing or pion mass) can be found in [26, 27].

### 2.2 Computation of the $Q\bar{Q}$ static potential in position space

First we determine the  $Q\bar{Q}$  static potential  $V(\vec{r})$  in position space. In a second step,  $V(\vec{r})$  is transformed to momentum space by means of a discrete Fourier transform.

In order to perform this Fourier transform, we need the static potential for all  $\vec{r} = \vec{n}a$  inside a periodic spatial volume  $L^3$  of side length  $L'$ . This is achieved as follows: First, we compute  $V(\vec{n}a)$  with  $n_i \in \{0, 1, \dots, N'/2\}$ , where  $N'$  is even and defined as  $N' \equiv L'/a$ . Second, we

$\beta$	$a$ in fm	$(L/a)^3 \times T/a$	$m_{\text{PS}}$ in MeV	$r_0/a$	# gauges
3.90	0.079(3)	$24^3 \times 48$	340(13)	5.36(4)	168
4.05	0.063(2)	$32^3 \times 64$	325(10)	6.73(5)	71
4.20	0.0514(8)	$48^3 \times 96$	284(5)	8.36(6)	46
4.35	0.0420(17)	$32^3 \times 64$	352(22)	9.81(13)	146

Table 1: Ensembles of gauge link configurations employed in the present study.

realize the periodicity by defining  $V(n_x a, n_y a, n_z a) \equiv V(|n_x|a, |n_y|a, |n_z|a)$ , where now  $n_i \in \{-\frac{N'}{2} + 1, -\frac{N'}{2} + 2, \dots, \frac{N'}{2}\}$ .

To keep finite volume effects on a negligible level, the spatial volume  $L^3$  needs to be sufficiently large, e.g. for our finest lattice spacing  $a \approx 0.0420$  fm we use  $L'/a = 256$ . While a lattice computation of the static potential  $V(\vec{n}a)$  for all  $n_x, n_y, n_z = 0, 1, \dots, N'/2$  is possible in principle, it is extremely computer time consuming in practice: One needs to generate gauge link configurations for such large volumes and has to compute both on- and off-axis Wilson loops for all possible quark-antiquark separations  $\vec{r} = \vec{n}a$ . On the other hand the shape of the static potential at large separations is known to be accurately described by  $V(r) = A_0 + \sigma r + A_1/r$ , where  $A_0$  denotes a constant offset parameter,  $\sigma$  is the string tension and  $A_1 \gtrsim -\pi/12$  [33–35]. Additionally, the large distance behavior of  $V(r)$  is expected to have a rather weak effect on the Fourier transformed potential for  $p \gg \Lambda_{\text{QCD}}$ , i.e. the momentum regime used for the matching to perturbation theory and the  $\Lambda_{\overline{\text{MS}}}$  determination in this work. Therefore, we stick to the following strategy:

(1) **Perform a standard lattice computation of  $V(\vec{r})$  for  $|\vec{r}| = |\vec{n}|a \leq r_{\text{max}}$ :**

For quark-antiquark separations  $\vec{n}a \leq r_{\text{max}}$  we extract  $V(\vec{r})$  from the exponential decay of Wilson loop averages  $\langle W(\vec{r}, t) \rangle$  with respect to their temporal extent  $t$ , while keeping their spatial extent  $r$  fixed. To this end we first compute

$$V^{(\text{effective})}(\vec{r}, t) = \frac{1}{a} \ln \left( \frac{\langle W(\vec{r}, t) \rangle}{\langle W(\vec{r}, t+a) \rangle} \right). \quad (4)$$

Somewhat arbitrarily, we choose  $r_{\text{max}} \approx 0.42$  fm corresponding to  $r_{\text{max}} = 10a$  for our smallest lattice spacing (cf. Table 1). In a second step the  $t$ -independent quantity  $V(\vec{r})$  is obtained by performing an uncorrelated  $\chi^2$  minimizing fit to  $V^{(\text{effective})}(\vec{r}, t)$  in a suitable  $t$  range. This range is chosen such that excited states are strongly suppressed, while statistical errors are still small.

We use the ensembles listed in Table 1, where  $L \ll L'$ , and consider on- and off-axis Wilson loops formed by APE smeared spatial links ( $N_{\text{APE}} = 60$ ,  $\alpha_{\text{APE}} = 0.5$  for all our gauge link ensembles) and ordinary, i.e. unsmeared temporal links. For a detailed explanation regarding the construction of off-axis Wilson loops cf. [5]. For a definition of APE smearing we refer to [36].

(2) **Model  $V(\vec{r})$  for  $|\vec{r}| = |\vec{n}|a > r_{\text{max}}$ :**

For quark-antiquark separations  $|\vec{n}|a > r_{\max}$  we model the lattice potential by

$$V(\vec{r}) = V_M(r) \equiv A_0 + \sigma r + \sum_{m=1}^M \frac{A_m}{r^m}. \quad (5)$$

In Sec. 4 different values of  $M \in \{1, 2, 3, 4\}$  are used to quantify systematic errors associated with this modeling of the long range part of the lattice potential. The string tension is fixed to  $\sigma = 1550 \text{ MeV/fm}$  (corresponding to  $r_0 = 0.420 \text{ fm}$  determined on the same gauge link configurations we are using throughout this work [27]). While  $A_1 = -\pi/12$  in the bosonic string picture [33, 34], lattice simulations with  $n_f = 2$  quark flavors yield a larger value  $A_1 \approx -0.3 \dots -0.5$  [35]. We determine  $A_m$  with  $m \in \{0, 1, \dots, M\}$  by a  $\chi^2$  minimizing fit of Eq. (5) to the lattice results determined in step (1) in the region  $r_{\min} \leq r \leq r_{\max}$ . In order to have  $\chi^2 \lesssim 1$ , for our smallest lattice spacing  $a = 0.0420 \text{ fm}$  we choose  $r_{\min} = 8a$  for  $M = 1$ ,  $r_{\min} = 6a$  for  $M = 2$ , and  $r_{\min} = 4a$  for  $M = \{3, 4\}$ .

The resulting function  $V_3(r)$  for  $M = 3$  is shown in Figure 1 together with the lattice results for  $V(\vec{r})$ .




Figure 1: Position space potential  $V_3(r)$  (green curve) obtained by a  $\chi^2$  minimizing fit to the lattice results  $V(\vec{r})$  (red dots) in the region  $4a \leq |\vec{r}| \leq 10a$  with  $a \approx 0.0420 \text{ fm}$ .

### 2.3 Computation of the $Q\bar{Q}$ static potential in momentum space

We define the  $Q\bar{Q}$  static potential in momentum space,  $V(\vec{p})$  with  $\vec{p} = 2\pi\vec{k}/L'$ , by the discrete Fourier transform of  $V(\vec{r})$ ,

$$V(\vec{p}) = V(2\pi\vec{k}/L') = \sum_{n_x, n_y, n_z} a^3 \exp\left(-\frac{2\pi i \vec{k} \vec{n}}{N'}\right) V(\vec{n}a), \quad (6)$$

where the sum is also over all possible values of  $n_i \equiv -\frac{N'}{2} + \nu_i$  with  $\nu_i \in \{1, 2, \dots, N'\}$ , and  $k_i \equiv -\frac{N'}{2} + \kappa_i$  with  $\kappa_i \in \{1, 2, \dots, N'\}$ .

In the continuum and in infinite volume  $V(\vec{p})$  is rotationally symmetric, i.e.  $V(\vec{p}) = V(p)$ . Since both a cubic lattice discretization and a cubic periodic volume break rotational symmetry, this is no longer true for the lattice potential (6). The deviations from the rotationally invariant continuum and infinite volume potential are expected to be particularly small, when restricting the dimensionless lattice momenta  $\vec{k}$  to values inside a cylinder of unit radius around the lattice diagonal, i.e. demanding

$$\vec{k}^2 - (\vec{k}\vec{d})^2 \leq 1, \quad \vec{d} = \frac{1}{\sqrt{3}}(1, 1, 1). \quad (7)$$

This so-called cylinder cut is frequently used in lattice momentum space computations of propagators. For details cf. [37].

The maximal momentum value along each of the three axes is  $\bar{p} = \pi/a \approx 15$  GeV for our smallest lattice spacing  $a \approx 0.0420$  fm. Similar to position space, where the minimum on-axis separation is  $a$  and the static potential is essentially free of discretization errors for  $r \gtrsim 3a$ , one might expect rather small discretization errors for  $p \lesssim \bar{p}/3 \approx 5$  GeV.

The final result  $V(p)$  for our smallest lattice spacing  $a \approx 0.0420$  fm, obtained with  $V_3(r)$  [cf. Eq. (5) and Figure 1] is shown in Figure 2<sup>3</sup>.




Figure 2:  $V(p)$  for  $a \approx 0.0420$  fm obtained from  $V_3(r)$  after applying the cylinder cut.

---

<sup>3</sup>Throughout this paper all computations are performed in units of the lattice spacing  $a$ , e.g. we work with quantities  $aV(r)$ ,  $V(p)/a^2$  and  $a\Lambda_{\overline{\text{MS}}}$ , which are independent of any potential errors or uncertainties regarding scale setting; for a recent review cf. [38]. Nevertheless, the axes in the presented plots as well as the numbers quoted in the main text often given in units of MeV or fm. This is intended to make the physical scales more obvious. To this end we use  $a = 0.0420$  fm for our smallest lattice spacing corresponding to  $\beta = 4.35$  (cf. also Table 1).



### 3 Perturbative insights into the $Q\bar{Q}$ static potential

#### 3.1 The $Q\bar{Q}$ static potential in perturbation theory

To allow for easier reference and to keep this paper self-contained, we briefly summarize the present status of the  $Q\bar{Q}$  static potential in perturbation theory. Quantities which depend on the particular renormalization scheme used are given in the  $\overline{\text{MS}}$  scheme [39, 40].

In the perturbative momentum regime, i.e. for momenta  $p = |\vec{p}| \gg \Lambda_{\text{QCD}}$ , the  $Q\bar{Q}$  static potential is conveniently represented as

$$V(p) = -C_F \frac{4\pi}{p^2} \alpha_V[\alpha_s(\mu), L(\mu, p)] \quad (8)$$

with  $C_F$  the eigenvalue of the quadratic Casimir operator for the fundamental representation of the gauge group;  $C_F = 4/3$  for  $\text{SU}(3)$ . Pulling out the overall factor  $\sim 1/p^2$  the entire non-trivial structure of  $V(p)$  can be encoded in the dimensionless quantity  $\alpha_V$ , which in turn is a function of both the coupling  $\alpha_s(\mu)$  and  $L \equiv L(\mu, p) = \ln \frac{\mu^2}{p^2}$ .  $\alpha_s(\mu)$  is evaluated at an *a priori* arbitrarily chosen renormalization scale  $\mu$  in the perturbative regime, i.e.  $\mu \gg \Lambda_{\text{QCD}}$ , guaranteeing that  $\alpha_s(\mu) \ll 1$ .

The running of the coupling  $\alpha_s(\mu)$  as a function of the renormalization scale  $\mu$  is governed by the QCD  $\beta$ -function defined as

$$\beta[\alpha_s(\mu)] \equiv \frac{\mu}{\alpha_s(\mu)} \frac{d}{d\mu} \alpha_s(\mu), \quad (9)$$

whose series expansion in powers of  $\alpha_s$  is presently known with the following accuracy,

$$\beta(\alpha_s) = -\frac{\alpha_s}{2\pi} \beta_0 \left[ 1 + \frac{\alpha_s}{4\pi} \frac{\beta_1}{\beta_0} + \left( \frac{\alpha_s}{4\pi} \right)^2 \frac{\beta_2}{\beta_0} + \left( \frac{\alpha_s}{4\pi} \right)^3 \frac{\beta_3}{\beta_0} + \dots \right]. \quad (10)$$

While the expansion coefficients  $\beta_0$  and  $\beta_1$  are independent of the renormalization scheme,  $\beta_2$  and  $\beta_3$  are scheme-dependent. They have been determined for arbitrary compact semi-simple Lie groups in the  $\overline{\text{MS}}$  scheme [41]. For  $\text{SU}(3)$  with  $n_f = 2$  massless dynamical quark flavors they read

$$\beta_0 = \frac{29}{3}, \quad \beta_1 = \frac{230}{3}, \quad \beta_2 = \frac{48241}{54}, \quad \beta_3 = \frac{18799309}{1458} + \frac{275524}{81} \zeta(3). \quad (11)$$

The same quantities for arbitrary values of  $n_f$  can be found e.g. in [5].

As the static potential is a physical observable, it should of course be independent of the explicit value of the renormalization scale  $\mu$  and form a renormalization group (RG) invariant, i.e. fulfill

$$\mu \frac{d}{d\mu} V(p) = 0. \quad (12)$$

One might wonder, how this can come about with  $\alpha_V$  in Eq. (8) being a function of the two  $\mu$ -dependent quantities  $\alpha_s(\mu)$  and  $L(\mu, p)$ . However, knowing  $V(p)$  – and thus  $\alpha_V$  – at a certain accuracy in perturbation theory, e.g. up to  $\mathcal{O}(\alpha_s^{\bar{k}})$ , Eq. (12) only has to hold to this order, i.e.

$$\mu \frac{d}{d\mu} \alpha_V[\alpha_s, L] = \mathcal{O}(\alpha_s^{\bar{k}+1}) \quad \leftrightarrow \quad \left( \frac{\partial}{\partial L} + \frac{\alpha_s}{2} \beta(\alpha_s) \frac{\partial}{\partial \alpha_s} \right) \alpha_V[\alpha_s, L] = \mathcal{O}(\alpha_s^{\bar{k}+1}). \quad (13)$$

Presently, all terms are known explicitly for  $\bar{k} = 4$ , and  $\alpha_V$  is of the following form,

$$\alpha_V[\alpha_s(\mu), L(\mu, p)] = \alpha_s(\mu) \left\{ 1 + \frac{\alpha_s(\mu)}{4\pi} P_1(L) + \left( \frac{\alpha_s(\mu)}{4\pi} \right)^2 P_2(L) + \left( \frac{\alpha_s(\mu)}{4\pi} \right)^3 \left[ P_3(L) + a_{3\ln} \ln \alpha_s(\mu) \right] + \dots \right\}. \quad (14)$$

While  $\alpha_V$  has a strict power-series expansion in  $\alpha_s$  up to  $\mathcal{O}(\alpha_s^3)$ , beyond this order one also encounters logarithmic contributions in  $\alpha_s$  [42], the first such term being  $\sim \alpha_s^4 \ln \alpha_s$  [43].

Equation (13) constrains the  $P_k(L)$  accounting for the entire  $L$  dependence of  $\alpha_V$  in Eq. (14) to be polynomials in  $L$  of degree  $k$ , i.e.

$$P_k(L) = \sum_{m=0}^k \rho_{km} L^m \quad (15)$$

with dimensionless expansion coefficients  $\rho_{km}$ , and implies that, apart from the explicit values of  $a_k \equiv \rho_{k0} = P_k(0)$  with  $a_0 = 1$ , the  $\rho_{km}$  are fully determined by the coefficients of the  $\beta$ -function [23, 44]. Their explicit expressions for  $k \leq 3$  are given in our notations in [23].

The coefficients  $a_1$  [19, 45] and  $a_2$  [46–48] are known analytically. For gauge group SU(3),  $n_f = 2$  and in the  $\overline{\text{MS}}$ -scheme they read

$$a_1 = \frac{73}{9}, \quad a_2 = \frac{25139}{162} + 9\pi^2 \left( 4 - \frac{\pi^2}{4} \right) + \frac{94}{3} \zeta(3). \quad (16)$$

Also the coefficients  $a_3$  and  $a_{3\ln}$  are known [43, 49–53]. Specializing to SU(3) and  $n_f = 2$ , they are given by (cf. [5])

$$a_3 = 27c_1 + \frac{15}{16}c_2 + 9c_3 + \frac{5}{48}c_4 - \frac{968981}{729} - 8\pi^2 \left( 15 - \frac{8\pi^2}{45} \right) + 144\pi^2 \left( \ln 3 + \gamma_E \right) + \frac{38192}{27} \zeta(3) + \frac{320}{9} \zeta(5) \quad (17)$$

and  $a_{3\ln} = 144\pi^2$ . The constants  $c_i$  ( $i = 1 \dots 4$ ) are only known numerically:

$$c_1 = 502.24(1), \quad c_2 = -136.39(12), \quad c_3 = -709.717, \quad c_4 = -56.83(1). \quad (18)$$

$c_1$  and  $c_2$  have been determined independently by both [50] and [51]. We use the numerical values from [50], who provide smaller statistical errors.<sup>4</sup>  $c_3$  and  $c_4$  have been determined by [49]. The coefficients  $a_i$  ( $i = 1 \dots 4$ ) for arbitrary values of  $n_f$  can be found, e.g., in [5].

Therewith, all coefficients in the perturbative expansion of the static potential in momentum space up to order  $\alpha_s^4$  have been assembled. We emphasize again that the resulting expression is independent of the explicit choice for  $\mu$ , in the sense that different choices for  $\mu$  only lead to deviations at  $\mathcal{O}(\alpha_s^5)$  [cf. Eq. (13)], which is beyond the accuracy of the contributions taken into account in Eq. (14).

---

<sup>4</sup>The errors associated with  $c_i$  ( $i = 1 \dots 4$ ) turn out to be negligible in the context of our  $\Lambda_{\overline{\text{MS}}}$  determination; therefore, we will not discuss them any further.

In order to prevent the logarithms  $L$  in Eq. (14) from becoming large and thereby spoil the perturbative expansion, it is desirable to ensure that  $\mu$  does not deviate much from  $p$ . Hence, a particularly convenient choice for the renormalization scale is  $\mu \equiv p$ , implying  $L = 0$ . We will exclusively stick to this choice throughout the remainder of this paper. Adopting this choice,  $\alpha_V$  simplifies significantly. It becomes a function of  $\alpha_s(p)$  only and reads

$$\alpha_V[\alpha_s(p)] = \alpha_s(p) \left\{ 1 + \frac{\alpha_s(p)}{4\pi} a_1 + \left( \frac{\alpha_s(p)}{4\pi} \right)^2 a_2 + \left( \frac{\alpha_s(p)}{4\pi} \right)^3 \left[ a_3 + a_{3\ln} \ln \alpha_s(p) \right] + \dots \right\}. \quad (19)$$

For completeness, note that – as briefly mentioned in the introduction – an analogous choice in position space, i.e. setting  $\mu \equiv 1/r$  completely spoils the convergence of the perturbative potential  $V(r)$  [19]. As will also become obvious below, the choice  $\mu \equiv p$  does not lead to any problems, the reason being the much less pronounced IR sensitivity of the perturbative potential  $V(p)$  in momentum space [11].

An alternative representation of Eq. (19) is

$$\frac{\alpha_V[\alpha_s(p)]}{\alpha_s(p)} = 1 + x a_1 + (x a_1)^2 \frac{a_2}{a_1^2} + (x a_1)^3 \frac{a_3 + a_{3\ln} \ln(4\pi/a_1)}{a_1^3} \left[ 1 + \frac{a_{3\ln}}{a_3 + a_{3\ln} \ln(4\pi/a_1)} \ln(x a_1) \right] + \dots, \quad (20)$$

where we employed the shortcut notation  $x \equiv \frac{\alpha_s(p)}{4\pi}$ .

In order to allow for more insights into the structure of  $\alpha_V[\alpha_s(p)]$  for  $n_f = 2$ , we insert the explicit numerical values for the coefficients  $a_1$ ,  $a_2$ ,  $a_3$  and  $a_{3\ln}$  into Eq. (20), and rewrite it as

$$\frac{\alpha_V[\alpha_s(p)]}{\alpha_s(p)} \approx 1 + x a_1 + 5.00 (x a_1)^2 + 17.63 (x a_1)^3 \left[ 1 + 0.15 \ln(x a_1) \right] + \dots, \quad (21)$$

with  $a_1 \approx 8.11$ . It is easy to check that for  $x a_1 \lesssim \frac{1}{5} \leftrightarrow \alpha_s(p) \lesssim \frac{4\pi}{5a_1} \approx 0.31$  the contributions  $\sim (x a_1)^n$  in Eq. (21) are ordered in the sense that they become increasingly less important, when increasing  $n$  from 0 to 3. For larger values of  $\alpha_s(p) \gtrsim 0.31$  this ordering is spoiled. Correspondingly, the perturbative expansion of  $\alpha_V[\alpha_s(p)]$  with  $n_f = 2$  can in particular be considered as well-behaved and controllable for  $\alpha_s(p) \ll 0.31$ . As the highest order contribution  $\sim (x a_1)^3$  in Eq. (21) is still significantly smaller than the next-to-highest one  $\sim (x a_1)^2$ , in the present paper we will consider the perturbative expansion as trustworthy even up to  $\alpha_s(p) \lesssim 0.31$ . Taking into account the value of  $\Lambda_{\overline{\text{MS}}}$  as determined in [5],  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 315 \text{ MeV}$ , this directly translates into a restriction to momenta  $p \gtrsim 1500 \text{ MeV}$  (cf. Sec. 3.2).

### 3.2 The scale $\Lambda_{\overline{\text{MS}}}$

So far we have not discussed, how an explicit numerical value can be attributed to the strong coupling  $\alpha_s(\mu)$  evaluated at a given momentum  $\mu$ . We emphasize that such an identification is renormalization scheme dependent, the most widely used scheme being the  $\overline{\text{MS}}$  scheme [39, 40], which we also adopt here.

A straightforward integration of Eq. (9) yields

$$\ln \frac{\mu}{\Lambda} = \left[ \int \frac{d\alpha_s}{\alpha_s} \frac{1}{\beta(\alpha_s)} \right] \Big|_{\alpha_s=\alpha_s(\mu)} + C, \quad (22)$$

where the conventional definition of  $\Lambda$  in the  $\overline{\text{MS}}$  scheme, i.e.  $\Lambda \rightarrow \Lambda_{\overline{\text{MS}}}$ , corresponds to the choice of  $C = \frac{\beta_1}{2\beta_0^2} \ln(\frac{\beta_0}{4\pi})$  [54]<sup>5</sup>.

Employing some elementary manipulations and rearrangements, Eq. (22) adopted to the  $\overline{\text{MS}}$  scheme can be written as

$$\Lambda_{\overline{\text{MS}}} \equiv \mu \left( \frac{\beta_0 \alpha_s(\mu)}{4\pi} \right)^{-\frac{\beta_1}{2\beta_0^2}} \exp \left\{ -\frac{2\pi}{\beta_0 \alpha_s(\mu)} - \int_0^{\alpha_s(\mu)} \frac{d\alpha_s}{\alpha_s} \left[ \frac{1}{\beta(\alpha_s)} + \frac{2\pi}{\beta_0 \alpha_s} - \frac{\beta_1}{2\beta_0^2} \right] \right\}, \quad (23)$$

involving a definite integral over  $\alpha_s$  (cf. e.g. [55]). The additional terms apart from the factor of  $1/\beta(\alpha_s)$  have been included in the integrand to make the finiteness of the integral over the interval from 0 to  $\alpha_s(\mu)$  manifest.

In general Eq. (23) cannot be solved explicitly to provide the coupling  $\alpha_s(\mu)$  at a given momentum scale  $\mu$  as a function of the ratio  $\mu/\Lambda_{\overline{\text{MS}}}$ . An exact closed form solution is only possible at leading order – i.e. taking into account only the leading contribution of the  $\beta$ -function (10),  $\beta(\alpha_s) \approx -\frac{\alpha_s}{2\pi}\beta_0$  – and reads  $\alpha_s(\mu) = \left[ \frac{\beta_0}{2\pi} \ln(\mu/\Lambda) \right]^{-1}$ . Note, however, that approximate results for  $\alpha_s(\mu)$  as a function of the ratio  $\mu/\Lambda_{\overline{\text{MS}}}$  are available also for higher order contributions: the derivation of such expressions involves expansions in terms of  $\ln(\mu/\Lambda_{\overline{\text{MS}}})$ ; see e.g. [54, 56].

Aiming at the determination of  $\Lambda_{\overline{\text{MS}}}$  by fitting the perturbative expression for the static potential (8) to numerical data from lattice simulations we do not see any reason to resort to these further approximations. Our strategy is rather to solve the implicit equation (23) for  $\alpha_s(\mu)$  numerically. Hence, we will always use the relation (23) between  $\alpha_s(\mu)$  and  $\mu/\Lambda_{\overline{\text{MS}}}$  at the presently best available accuracy, irrespectively of the order of the expansion of the perturbative potential in  $\alpha_s(p)$ .

However, note that there is still some freedom left in deciding, how to proceed with the evaluation of Eq. (23). We can

- (I) either plug in the perturbative expression of the  $\beta$ -function (10) at the presently best known accuracy and then do the integration over  $\alpha_s$  numerically,
- (II) or adopt a Taylor expansion of the integrand in Eq. (23) and do the integral analytically, keeping only those terms, whose coefficients are known explicitly,

$$\begin{aligned} \int_0^{\alpha_s(\mu)} \frac{d\alpha_s}{\alpha_s} \left[ \frac{2}{\beta(\alpha_s)} + \frac{4\pi}{\beta_0 \alpha_s} - \frac{\beta_1}{\beta_0^2} \right] \\ = \frac{\beta_0 \beta_2 - \beta_1^2}{\beta_0^3} \frac{\alpha_s(\mu)}{4\pi} + \frac{\beta_0^2 \beta_3 - 2\beta_0 \beta_1 \beta_2 + \beta_1^3}{2\beta_0^4} \left( \frac{\alpha_s(\mu)}{4\pi} \right)^2 + \mathcal{O}(\alpha_s^3). \end{aligned} \quad (24)$$

Of course, limiting ourselves to the truly perturbative regime, i.e. the regime, where the perturbative expansion of the  $\beta$ -function (10) is such that higher order corrections become increasingly less important, both choices should be equally justified.

As pointed out in Sec. 3, when fitting the perturbative static potential (8) in momentum space to lattice data we will always stick to the identification  $\mu \equiv p$ . Correspondingly, higher order

---

<sup>5</sup>For completeness, note that our conventions slightly differ from those of [54], who write Eq. (9) in terms of  $a \equiv \alpha_s/\pi$ . Moreover,  $\beta_n|_{[54]} = \beta_n/4^{n+1}$  and  $b_n|_{[54]} = \beta_n/(4^n \beta_0)$ .

corrections in Eq. (10) should remain small throughout the integration interval from 0 to  $\alpha(p)$  in Eq. (23) and Eq. (24), respectively.

To get a feeling, how the perturbative expansion of the  $\beta$ -function behaves, we employ the same strategy as adopted in the context of Eq. (21) and study its behavior at the upper integration limit in Eq. (23), i.e. at  $\alpha_s \equiv \alpha_s(p)$ . First we rewrite Eq. (10) as

$$\beta(\alpha_s) = -\frac{\alpha_s}{2\pi}\beta_0 \left[ 1 + x \frac{\beta_1}{\beta_0} + \left( x \frac{\beta_1}{\beta_0} \right)^2 \frac{\beta_0\beta_2}{\beta_1^2} + \left( x \frac{\beta_1}{\beta_0} \right)^3 \frac{\beta_0^2\beta_3}{\beta_1^3} + \dots \right] \quad (25)$$

with  $x \equiv \frac{\alpha_s(p)}{4\pi}$ . Second we insert the explicit numerical coefficients of the  $\beta$ -function for  $n_f = 2$  into this equation, resulting in

$$\beta(\alpha_s) \approx -\frac{\alpha_s}{2\pi}\beta_0 \left[ 1 + x \frac{\beta_1}{\beta_0} + 1.47 \left( x \frac{\beta_1}{\beta_0} \right)^2 + 3.52 \left( x \frac{\beta_1}{\beta_0} \right)^3 + \dots \right] \quad (26)$$

with  $\frac{\beta_1}{\beta_0} \approx 7.93$ . Using the same reasoning as in the context of Eq. (21) above, we find that for the contributions  $\sim (x\beta_1/\beta_0)^n$  to become increasingly less important with  $n$ , we have to demand  $x\beta_1/\beta_0 \lesssim 0.4$ , which corresponds to  $\alpha_s(p) \lesssim 0.63$ .

Hence, in particular for  $\alpha_s(p) \lesssim 0.31$  – which was the value found to crudely delimit the range of validity of a perturbative expansion of the static potential in momentum space for  $n_f = 2$  [cf. in the context of Eq. (21) above] – higher order terms in the perturbative expansion of the  $\beta$ -function (10) for  $n_f = 2$  are expected to become much less important. In turn both possibilities (I) and (II) discussed above to numerically solve Eq. (23) for  $\alpha_s(p)$  as a function of  $p/\Lambda_{\overline{\text{MS}}}$  should yield very similar results.

## 4 Determination of $\Lambda_{\overline{\text{MS}}}$

In this section, we determine  $\Lambda_{\overline{\text{MS}}}$  in units of the lattice spacing, i.e.  $a\Lambda_{\overline{\text{MS}}}$ , by fitting perturbative expressions for the static potential in momentum space (cf. Sec. 3) to corresponding lattice results (cf. Sec. 2). Using the values of the lattice spacing listed in Table 1, these results can easily be converted to physical units, i.e. MeV. Unless explicitly stated otherwise, the errors provided for  $\Lambda_{\overline{\text{MS}}}$  do not account for the uncertainties associated with the lattice spacing  $a$ . These errors will however be accounted for when quoting our final result for  $\Lambda_{\overline{\text{MS}}}$  [cf. Eq. (31), below]. For completeness, we will also quote  $r_0\Lambda_{\overline{\text{MS}}}$ ,<sup>6</sup> which is dimensionless and, hence, unaffected by any potential uncertainty in  $a$  (uncertainties in  $r_0/a$ , which are collected in Table 1, are, of course, included).

### 4.1 Fitting procedures

The perturbative  $Q\bar{Q}$  potential to be fitted to lattice results is given by

$$V(p) = -\frac{4}{3} \frac{4\pi}{p^2} \alpha_s(p) \left\{ \underbrace{1}_{\text{LO}} + \underbrace{\frac{\alpha_s(p)}{4\pi} a_1}_{\text{NLO}} + \underbrace{\left(\frac{\alpha_s(p)}{4\pi}\right)^2 a_2}_{\text{NNLO}} + \underbrace{\left(\frac{\alpha_s(p)}{4\pi}\right)^3 [a_3 + a_{3\ln} \ln \alpha_s(p)]}_{\text{NNNLO}} \right\} + V_0, \quad (27)$$

where we included an overall constant offset  $V_0$  of the potential. We use different orders in the expansion of  $V(p)$  in powers of  $\alpha_s(p)$  to test and judge the convergence behavior of our results; the abbreviation N<sup>n</sup>LO stands for (next-to-)<sup>n</sup>leading-order. As only energy differences are measurable  $V_0$  does not have any observable consequences. However, it is necessary to allow for a meaningful matching of the perturbative potential to lattice results.

For the relation between  $\alpha_s(p)$  and  $\Lambda_{\overline{\text{MS}}}$  we employ (cf. Sec. 3)

- (I) either Eq. (23) with  $\beta(\alpha)$  given by the terms written explicitly in Eq. (10),
- (II) or Eq. (23) with the integral expression substituted for the terms written explicitly on the left-hand side of Eq. (24).

These implicit equations are solved numerically to yield  $\alpha_s(p)$  as a function of  $p/\Lambda_{\overline{\text{MS}}}$ .

We employ an uncorrelated  $\chi^2$  minimizing fit with two degrees of freedom,  $V_0$  and  $\Lambda_{\overline{\text{MS}}}$ , to fit the perturbative  $Q\bar{Q}$  potential (27) to the lattice potential in momentum space. Our fitting interval is delimited by  $p_{\min}$  and  $p_{\max}$ . Note that the lattice potential in momentum space originates from position space results whose large distance behavior is modeled by Eq. (5) with  $M \in \{1, 2, 3, 4\}$  – and thus also depends on  $M$  (cf. Sec. 2).

---

<sup>6</sup>The scale  $r_0$  is defined via  $r_0^2 F(r_0) = 1.65$ , with  $F(r) = dV(r)/dr$  [57].

## 4.2 Systematic and statistical errors of $\Lambda_{\overline{\text{MS}}}$

### 4.2.1 Individual variation of input parameters

We investigate the stability of  $\Lambda_{\overline{\text{MS}}}$  with respect to variations of the input parameters  $p_{\min}$ ,  $p_{\max}$  and  $M$ , using our finest lattice spacing  $a \approx 0.042$  fm and  $(L'/a)^3 = 256^3$  lattice sites. To this end the input parameters delimiting the fitting range are varied in the following intervals:

- $p_{\min} = 1500$  MeV  $\dots$  2250 MeV:  
As discussed in detail in Sec. 3, for  $p_{\min} \lesssim 1500$  MeV the validity of perturbative expressions for the static potential is rather questionable.
- $p_{\max} = 2250$  MeV  $\dots$  3000 MeV:  
The maximum momentum on our finest lattice along an axis is  $\bar{p} = \pi/a \approx 15$  GeV. For  $p \lesssim \bar{p}/3$ , it seems reasonable to expect rather small discretization errors. These expectations have also been confirmed by numerical investigations. Fitting the perturbative potential to lattice results we obtain an essentially stable value of  $\Lambda_{\overline{\text{MS}}}$  up to 3000 MeV  $\dots$  3500 MeV.

The different choices for the parameter  $M$  are

- $M \in \{1, 2, 3, 4\}$ :  
The parameter  $M$  is varied to estimate the systematic errors associated with our modeling of the long range part of the lattice potential in position space; cf. Sec. 2.2.

Below, we demonstrate that the fit results for  $\Lambda_{\overline{\text{MS}}}$  are rather stable with respect to these parameter variations, i.e. we confirm that a meaningful and rather precise matching of perturbative expressions for the  $Q\bar{Q}$  static potential and lattice results is possible.

Exemplary fits of the perturbative static potential (27) at LO, NLO, NNLO and NNNLO to lattice results are depicted in Figure 3. Noteworthy, all four orders are suited to describe the lattice potential within statistical errors, i.e. fulfill  $\chi^2/\text{dof} \lesssim 1.0$ . Similar  $\chi^2/\text{dof}$  are obtained when varying input parameters (cf. below).

To understand, how the value of  $\Lambda_{\overline{\text{MS}}}$  depends on the input parameters  $p_{\min}$  and  $p_{\max}$ , we vary them individually. Our results are summarized graphically in Figure 4:<sup>7</sup>

- The plots in the left column are obtained with fitting procedure (I), while those in the right column result from fitting procedure (II).
- For the plots in the first line we vary  $p_{\min} = 1500$  MeV  $\dots$  2250 MeV, while keeping  $p_{\max} = 2625$  MeV fixed at the center of the interval defined above, and set  $M = 3$ . We find:
  - Variation of  $\Lambda_{\overline{\text{MS}}}$  (NNLO):  
337 MeV  $\dots$  344 MeV (fitting procedure (I)),  
335 MeV  $\dots$  342 MeV (fitting procedure (II)).

---

<sup>7</sup>Statistical errors associated with the  $\chi^2$  minimizing fits, which are rather small ( $\approx 2$  MeV  $\dots$  4 MeV), are not considered in this subsection. They are, however, included in the final results presented in Sec. 4.3.




Figure 3: Exemplary fits of the perturbative static potential (27) to lattice results obtained with  $\beta = 4.35$  and  $M = 3$ . We employ fitting procedure (I) with  $p_{\min} = 1875$  MeV and  $p_{\max} = 2625$  MeV. The errors quoted below the plots refer to the fitting and are statistical errors only; systematic errors will be discussed below.

- Variation of  $\Lambda_{\overline{\text{MS}}}$  (NNNLO):  
 313 MeV ... 317 MeV (fitting procedure (I)),  
 312 MeV ... 315 MeV (fitting procedure (II)).
- For the plots in the second line we vary  $p_{\max} = 2250$  MeV ... 3000 MeV, while keeping  $p_{\min} = 1875$  MeV fixed at the center of the interval defined above, and set  $M = 3$ . We obtain:
  - Variation of  $\Lambda_{\overline{\text{MS}}}$  (NNLO):  
 338 MeV ... 343 MeV (fitting procedure (I)),  
 337 MeV ... 341 MeV (fitting procedure (II)).
  - Variation of  $\Lambda_{\overline{\text{MS}}}$  (NNNLO):  
 314 MeV ... 317 MeV (fitting procedure (I)),  
 313 MeV ... 315 MeV (fitting procedure (II)).
- For the plots in the third line we vary the center of the fitting range  $(p_{\min} + p_{\max})/2 = 1875$  MeV ... 2625 MeV, while keeping the width of the fitting range  $p_{\max} - p_{\min} = 750$  MeV fixed, and set  $M = 3$ . This results in:
  - Variation of  $\Lambda_{\overline{\text{MS}}}$  (NNLO):  
 335 MeV ... 340 MeV (fitting procedure (I)),  
 334 MeV ... 338 MeV (fitting procedure (II)).



- Variation of  $\Lambda_{\overline{\text{MS}}}^{\text{(NNNLO)}}$ :
  - 313 MeV ... 315 MeV (fitting procedure (I)),
  - 312 MeV ... 314 MeV (fitting procedure (II)).

To allow for a meaningful determination of  $\Lambda_{\overline{\text{MS}}}$ , it is important to ensure that the fit results for  $\Lambda_{\overline{\text{MS}}}$  only exhibit a weak – preferably negligible – dependence on  $p_{\min}$  and  $p_{\max}$ . In plots of  $\Lambda_{\overline{\text{MS}}}$  as a function of  $p_{\min}$  and  $p_{\max}$  this should manifest itself in the formation of plateaus. This behavior is clearly visible in Figure 4 for all six plots and at all orders in the expansion (27). Moreover, the curves shown in Figure 4 clearly reveal a convergence behavior when increasing the order in Eq. (27) from LO to NNNLO: First,  $\Lambda_{\overline{\text{MS}}|\text{LO}} > \Lambda_{\overline{\text{MS}}|\text{NLO}} > \Lambda_{\overline{\text{MS}}|\text{NNLO}} > \Lambda_{\overline{\text{MS}}|\text{NNNLO}}$ , second the ratios of their relative differences scale as

$$\Lambda_{\overline{\text{MS}}|\text{LO}} - \Lambda_{\overline{\text{MS}}|\text{NLO}} : \Lambda_{\overline{\text{MS}}|\text{NLO}} - \Lambda_{\overline{\text{MS}}|\text{NNLO}} : \Lambda_{\overline{\text{MS}}|\text{NNLO}} - \Lambda_{\overline{\text{MS}}|\text{NNNLO}} \approx 5 : 3 : 1. \quad (28)$$

Third, the values of  $\Lambda_{\overline{\text{MS}}}$  extracted from fits of Eq. (27) at NNLO and NNNLO to lattice results yield quite similar results. This can be considered as indication that the neglected orders beyond NNNLO in Eq. (27) will not alter the value of  $\Lambda_{\overline{\text{MS}}}$  significantly. As an – rather conservative – estimate of the systematic error due to the truncation of the perturbative expansion (27) at  $\mathcal{O}(\alpha_s^4)$  we take the difference between the NNLO and the NNNLO results for  $\Lambda_{\overline{\text{MS}}}$  (cf. Sec. 4.2.2).

Another source of uncertainty in our analysis is the modeling (5) used to extrapolate the lattice potential computed from Wilson loops to larger  $r$ . In order to scrutinize this uncertainty, we study the dependence of  $\Lambda_{\overline{\text{MS}}}$  on  $M \in \{1, 2, 3, 4\}$ , resorting to fitting procedure (I) and keeping  $p_{\min} = 1875 \text{ MeV}$  and  $p_{\max} = 2625 \text{ MeV}$  fixed. The corresponding results, which are collected in Table 2, indicate that  $\Lambda_{\overline{\text{MS}}}$  is also rather stable with respect to variations of  $M$ .

	$V_1(r)$	$V_2(r)$	$V_3(r)$	$V_4(r)$
NNLO, fitting procedure (I)	349	343	340	346
NNLO, fitting procedure (II)	347	341	338	345
NNNLO, fitting procedure (I)	323	318	315	321
NNNLO, fitting procedure (II)	322	317	314	320


Table 2: Results for  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$  in MeV for different  $M \in \{1, 2, 3, 4\}$  using fitting procedure (I) and  $p_{\min} = 1875 \text{ MeV}$  and  $p_{\max} = 2625 \text{ MeV}$ .

The variation of  $\Lambda_{\overline{\text{MS}}}$  as a function of the input parameters  $p_{\min}$ ,  $p_{\max}$  and  $M$  (cf. also Figure 4 and Table 2) when fitting (27), both at NNLO (red) and at NNNLO (green), to lattice results is visualized in Figure 5. As systematic uncertainty one could e.g. quote the whole range of values covered by the NNNLO variations,


$$\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 314 \text{ MeV} \dots 323 \text{ MeV}. \quad (29)$$

In particular note the drastic improvement in stability as compared to the previous position space analysis [5], where a systematic uncertainty larger by a factor of  $\approx 4$  has been obtained,  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 304 \text{ MeV} \dots 344 \text{ MeV}$  (cf. Eq. (46) and Figure 5 of [5]).


The sources of the systematic error discussed above might, however, be correlated. A method to determine the overall systematic error accounting for potential correlation is discussed in the following subsection.




(a) fitting proc. (I),  $p_{\max} = 2625$  MeV




(b) fitting proc. (II),  $p_{\max} = 2625$  MeV




(c) fitting proc. (I),  $p_{\min} = 1875$  MeV



(d) fitting proc. (II),  $p_{\min} = 1875$  MeV



(e) fitting proc. (I),  $p_{\max} - p_{\min} = 750$  MeV



(f) fitting proc. (II),  $p_{\max} - p_{\min} = 750$  MeV

Figure 4: Results for  $\Lambda_{\overline{\text{MS}}}$  obtained by fitting Eq. (27) to lattice results using  $M = 3$ ; **left column:** fitting procedure (I); **right column:** fitting procedure (II); **first line:**  $\Lambda_{\overline{\text{MS}}}$  as a function of  $p_{\max}$ ; **second line:**  $\Lambda_{\overline{\text{MS}}}$  as a function of  $p_{\min}$ ; **third line:**  $\Lambda_{\overline{\text{MS}}}$  as a function of  $(p_{\min} + p_{\max})/2$ .




Figure 5: Graphical summary of the variation of  $\Lambda_{\overline{\text{MS}}}$ , when varying the input parameters  $p_{\min}$ ,  $p_{\max}$  and  $M$  (cf. also Figure 4, Table 2 and the corresponding paragraphs in the main text). For each line, we indicate the fitting procedure used, i.e., either (I) and (II), and specify which parameter is varied. The variation is over the intervals specified at the begin of Sec. 4.2.1. The other input parameters are held fixed at the center of the respective intervals, and  $M = 3$  unless otherwise stated.

#### 4.2.2 Consideration of correlations between different systematic error sources

To account for possible correlations, we perform a large number of fits, with the input parameters chosen randomly and uniformly in the intervals specified in Sec. 4.2.1. As systematic error we then take the variance of the fit results.

We have performed 40,000 fits (i.e., sufficiently many to render the statistical error of the variance negligible):

- 10,000 NNLO fits, fitting procedure (I);
- 10,000 NNLO fits, fitting procedure (II);
- 10,000 NNNLO fits, fitting procedure (I);
- 10,000 NNNLO fits, fitting procedure (II).

In these fits we randomly vary  $p_{\min}$  and  $p_{\max}$  in the intervals  $p_{\min} = 1500 \text{ MeV} \dots 2250 \text{ MeV}$  and  $p_{\max} = 2250 \text{ MeV} \dots 3000 \text{ MeV}$  while imposing the constraint  $p_{\max} - p_{\min} \geq 375 \text{ MeV}$ . Moreover, we cyclically vary  $M \in \{1, 2, 3, 4\}$ . From this analysis we obtain an average and a variance, i.e. a systematic error, of

$$\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 331(13) \text{ MeV}. \quad (30)$$

Again it is most instructive to compare the error with that obtained for a position space analysis of the  $Q\bar{Q}$  static potential based upon the the same lattice setup [5]:

- The final result of Ref. [5] is  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 315(26)$  MeV. Hence, the momentum space analysis pursued in the present paper yields a result which is more precise by a factor of  $\approx 2$ .
- While fitting procedure (A) used in [5] is rather similar to (I) and (II) of this paper, fitting procedure (B) used in [5] is somewhat different.<sup>8</sup> Hence, in order to allow for a consistent and fair comparison of our present momentum space analysis and the previous position space analysis, only fitting procedure (A) should be used. We carried out such an analysis, resulting in  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 331(20)$  MeV. While there is now perfect agreement with respect to the average, the error associated with a momentum space analysis is still smaller by a factor of  $\approx 1.5$  than the error obtained in a position space analysis.
- Finally, one could be less conservative with regard to the estimate of the uncertainty associated with the truncation of the perturbative expressions and only take into account the NNNLO results. A momentum space analysis would then result in  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 318(3)$  MeV, while a position space analysis (fitting procedure (A) only) would lead to  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 326(13)$  MeV. In this case the error of  $\Lambda_{\overline{\text{MS}}}$  as obtained from a momentum space analysis is even smaller by a factor of  $\approx 4$  as compared to an analogous position space analysis. This rather drastic difference can be attributed to the almost perfect stability of  $\Lambda_{\overline{\text{MS}}}$  at NNNLO with respect to variations of the fitting range in momentum space (cf. Figure 4). In position space, a similar stability has not been observed [5].

Let us emphasize again that all these results for  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$  agree within statistical errors.

As already mentioned in footnote 7, the fitting procedure introduces an additional statistical uncertainty of  $\approx 2$  MeV  $\dots$  4 MeV, which we add in quadrature when generating the final result (cf. Sec. 4.3).

Further systematic uncertainties, exclusively originating from the lattice static potential, are discussed in the next section.

### 4.2.3 Systematic errors of $\Lambda_{\overline{\text{MS}}}$ associated with the lattice computation

#### Lattice discretization errors

To estimate the order of magnitude of lattice discretization errors, we utilize the lattice ensembles listed in Table 1, featuring pions of roughly of the same mass  $\approx 284$  MeV  $\dots$  352 MeV. Keeping the input parameters  $p_{\min} = 1300$  MeV,  $p_{\max} = 2050$  MeV and  $M = 3$  fixed, we extract values for both  $r_0\Lambda_{\overline{\text{MS}}}$  and  $\Lambda_{\overline{\text{MS}}}$  by performing the corresponding fits. For these investigations we exclusively adopt fitting procedure (I) and use the perturbative  $Q\bar{Q}$  static potential (27) at the

---

<sup>8</sup>In fact, fitting procedure (B) of [5] is expected to increase the error because the one-loop result for  $\alpha_s$ , employed in this fitting procedure, deviates significantly from the higher-order expressions for  $\alpha_s$  throughout the considered fitting interval.

best known accuracy, i.e. NNNLO.<sup>9</sup> In Figure 6 we depict our results for  $r_0\Lambda_{\overline{\text{MS}}}$  and  $\Lambda_{\overline{\text{MS}}}$  as a function of  $a^2$ . The results determined at the four available lattice spacings perfectly agree within errors, i.e. there is no indication of any sizable lattice discretization errors.




Figure 6: Dependence of  $r_0\Lambda_{\overline{\text{MS}}}$  and  $\Lambda_{\overline{\text{MS}}}$  on the lattice spacing. The result of an continuum extrapolation is depicted in green.

For completeness note that the errors associated with the extracted values for  $\Lambda_{\overline{\text{MS}}}$  in Figure 6 are rather large. The reason for this is that we have here included the errors associated with the lattice spacings and also with  $r_0/a$  (cf. Table 1). This is absolutely essential when comparing results obtained at different values of the lattice spacing.

Figure 6 also shows continuum extrapolations assuming a dependence  $\propto a^2$  for both  $r_0\Lambda_{\overline{\text{MS}}}$  and  $\Lambda_{\overline{\text{MS}}}$ . Within errors the results of these extrapolations agree with the results obtained for our smallest lattice spacing ( $\beta = 4.35$ ,  $a \approx 0.042$  fm). Even though no clear indication of any systematic upward or downward tendency as a function of  $a^2$  is visible in Figure 6, we are conservative in our error estimate and infer additional systematic errors of  $\pm 8$  MeV for  $\Lambda_{\overline{\text{MS}}}$  and  $\pm 0.010$  for  $r_0\Lambda_{\overline{\text{MS}}}$  to be accounted in our final result. These estimates are obtained by taking the difference between the central values of  $r_0\Lambda_{\overline{\text{MS}}}$  and  $\Lambda_{\overline{\text{MS}}}$  at our smallest lattice spacing and the corresponding continuum extrapolations.

## Finite volume effects and non-vanishing light quark mass corrections

Finite volume effects were investigated in detail in Ref. [5]. In this analysis such effects were found to be negligible compared to the other errors discussed above.

Similarly, potential corrections on  $\Lambda_{\overline{\text{MS}}}$  due to non-vanishing light quark masses were examined in detail in [5] by studying the variation of  $\Lambda_{\overline{\text{MS}}}$  for different pion masses in the range  $m_{\text{PS}} \approx 325 \text{ MeV} \dots 517 \text{ MeV}$  at fixed lattice spacing and spacetime volume. In the quark mass region investigated,  $\Lambda_{\overline{\text{MS}}}$  was found to be constant within tiny statistical errors of  $\approx \pm 1$  MeV. Therefore, we do not expect the non-vanishing light quark masses on the lattice to induce any significant deviations from the zero quark mass limit, for which the perturbative expressions in Sec. 3 were derived. In other words, we consider the systematic error introduced by comparing a lattice computation featuring massive light quarks with a perturbative calculation in the zero quark

<sup>9</sup>To allow for stable fits with sufficiently many lattice points for  $V(p)$  inside the interval  $p_{\text{min}} \dots p_{\text{max}}$ , with  $p_{\text{max}}$  not too close to the maximum lattice momentum along one of the three spatial axes, we had to choose  $p_{\text{min}}$  slightly smaller than the minimum value of  $p_{\text{min}} = 1500$  MeV defined in Sec. 4.2.1.

mass limit negligible compared to the other uncertainties discussed above.

For the above reasons we do not take any potential errors arising from finite volume effects and due to non-vanishing light quark mass corrections into account when quoting our final result for  $\Lambda_{\overline{\text{MS}}}$  in this paper.

### 4.3 Final results for $\Lambda_{\overline{\text{MS}}}$

In the following, we present our final results for  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$ . We quote these results both in units of MeV and in units of  $r_0$ . Since there seem to be up to  $\mathcal{O}(10\%)$  unresolved differences regarding scale setting between different lattice QCD collaborations [38], we prefer the scale setting-independent quantity  $r_0\Lambda_{\overline{\text{MS}}}$ .

The determination is based on the lattice results at our finest lattice spacing, i.e. at  $\beta = 4.35$ , as explained in Sec. 4.2.2. The individual error contributions, which we combine by adding them in quadrature, are

- (1) the correlated systematic errors associated with the unknown contributions beyond  $\mathcal{O}(\alpha_s^4)$  of the perturbative  $Q\bar{Q}$  static potential and the input parameters of the fitting procedure (cf. Sec. 4.2.2),
- (2) the statistical errors associated with the  $\chi^2$  minimizing fits,
- (3) the estimated lattice discretization errors (cf. Sec. 4.2.3), and
- (4) the errors associated with the lattice spacing  $a = 0.0420(17)$  fm and with  $r_0/a = 9.81(13)$ .

Taking all these contributions into account, we finally obtain

$$r_0\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 0.692(31), \quad \Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 331(21) \text{ MeV}. \quad (31)$$

For completeness, note that being less conservative with regard to the estimate of the uncertainty associated with the truncation of the perturbative expressions and just taking NNNLO fits into account (cf. also Sec. 4.2.2), would result in  $r_0\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 0.665(16)$  and  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 318(16)$  MeV.

The final results (31) for  $\Lambda_{\overline{\text{MS}}}$  extracted from our momentum space analysis are more precise than those obtained in position space using the same lattice setup:  $r_0\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 0.658(55)$  and  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 315(30)$  MeV.

Let us emphasize again that the errors quoted here and in Eq. (31) also account for contributions which are not directly related to the extraction of  $\Lambda_{\overline{\text{MS}}}$ , as e.g. the uncertainties of  $a$  and of  $r_0/a$ . Hence, with regard to a comparison of the results for  $\Lambda_{\overline{\text{MS}}}$  obtained from an analysis of the  $Q\bar{Q}$  static potential in momentum space as opposed to a similar analysis in position space, we consider the comparison in Sec. 4.2.2 – not accounting for these additional uncertainties – as most relevant and meaningful.

## 5 Conclusions

We have determined  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$  by fitting perturbative expressions for the  $Q\bar{Q}$  static potential at the presently best known accuracy, i.e. up to  $\mathcal{O}(\alpha_s^4)$  in momentum space, to lattice results obtained at a rather fine lattice spacing  $a \approx 0.042$  fm.

In contrast to previous works in this direction (cf. e.g. [1–7]) we have employed a discrete Fourier transform to transform the lattice results for the  $Q\bar{Q}$  static potential to momentum space. The extraction of  $\Lambda_{\overline{\text{MS}}}$  by fitting perturbative expressions of the static potential to lattice results has exclusively been performed in momentum space.

Resorting to a previous position space extraction of  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$  based on the the same lattice data [5], we could show that the momentum space analysis allows for a more precise determination of  $\Lambda_{\overline{\text{MS}}}$ . We could reduce the associated errors by a factor of  $\approx 1.5 \dots 4$ , depending on the details of the fitting procedures and the estimate of the error associated with the truncation of the perturbative expressions used (cf. in particular Sec. 4.2.2). This improvement can mainly be attributed to the nearly perfect stability of the results for  $\Lambda_{\overline{\text{MS}}}$  with respect to variations of the momentum fitting range (cf. Figure 4). Such behavior has not been observed in position space, where comparably rather strong variations have been observed (cf. also Figures 3 and 4 of [5]).

In units of the hadronic scale  $r_0$  our final result for  $\Lambda_{\overline{\text{MS}}}$  reads

$$r_0 \Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 0.692(31), \quad (32)$$

while in physical units it is given by

$$\Lambda_{\overline{\text{MS}}}^{(n_f=2)} = 331(21) \text{ MeV}. \quad (33)$$

The quoted uncertainties include all sources of systematic error: the neglect of higher orders in the perturbative expansion, the dependence of the fit results on the fitting range  $p_{\min} \dots p_{\max}$  and the parameter  $M$ , lattice discretization errors, finite volume effects and uncertainties due to nonvanishing quark masses on the lattice. The uncertainties are dominated by the variations of  $\Lambda_{\overline{\text{MS}}}$ , when switching from NNLO to NNNLO, and in the case of Eq. (33) by the uncertainties associated with the lattice spacing.

We note that the results (32) and (33) compare well with other determinations of  $\Lambda_{\overline{\text{MS}}}^{(n_f=2)}$  from the literature, e.g. [2, 4, 58–62] (cf. also the more detailed discussion and graphical summary in [5]).

In the future it would be interesting to adopt a similar momentum space analysis to lattice results for the  $Q\bar{Q}$  static potential with  $n_f = 0$ ,  $n_f = 2 + 1$  and  $n_f = 2 + 1 + 1$  dynamical quark flavors. For all these cases we expect a momentum space analysis to benefit from the better convergence behavior of the perturbative static potential in momentum space as compared to position space (cf. also the detailed discussion in the introduction, Sec. 1), which – as demonstrated in the present work – can reduce the error on the extracted values of  $\Lambda_{\overline{\text{MS}}}$  as compared to a position space analysis of the  $Q\bar{Q}$  static potential.

## Acknowledgments

We would like to thank Karl Jansen for helpful discussions, and the European Twisted Mass Collaboration for the generation of gauge link configurations used in this paper.

M.W. acknowledges support by the Emmy Noether Programme of the DFG (German Research Foundation), grant WA 3000/1-1. F.K. is grateful to M. & M. Tingu for inspiring comments, and thankful to Karl-Heinz “Servūs” Dubansky for providing a clean environment and Gummibären.

This work was supported in part by the Helmholtz International Center for FAIR within the framework of the LOEWE program launched by the State of Hesse.

## References

- [1] C. Michael, “The Running coupling from lattice gauge theory,” *Phys. Lett. B* **283**, 103 (1992) [arXiv:hep-lat/9205010].
- [2] M. Gockeler, R. Horsley, A. C. Irving, D. Pleiter, P. E. L. Rakow, G. Schierholz and H. Stuben, “A Determination of the Lambda parameter from full lattice QCD,” *Phys. Rev. D* **73**, 014513 (2006) [hep-ph/0502212].
- [3] N. Brambilla, X. Garcia i Tormo, J. Soto and A. Vairo, “Precision determination of  $r_0\Lambda_{\overline{\text{MS}}}$  from the QCD static energy,” *Phys. Rev. Lett.* **105**, 212001 (2010) [Erratum-ibid. **108**, 269903 (2012)] [arXiv:1006.2066 [hep-ph]].
- [4] B. Leder *et al.* [ALPHA Collaboration], “Scale  $r_0$  and the static potential from the CLS lattices,” *PoS LATTICE* **2010**, 233 (2010) [arXiv:1012.1141 [hep-lat]].
- [5] K. Jansen *et al.* [ETM Collaboration], “ $\Lambda_{\overline{\text{MS}}}$  from the static potential for QCD with  $n_f = 2$  dynamical quark flavors,” *JHEP* **1201**, 025 (2012) [arXiv:1110.6859 [hep-ph]].
- [6] A. Bazavov, N. Brambilla, X. Garcia i Tormo, P. Petreczky, J. Soto and A. Vairo, “Determination of  $\alpha_s$  from the QCD static energy,” *Phys. Rev. D* **86**, 114031 (2012) [arXiv:1205.6155 [hep-ph]].
- [7] P. Fritzscht, F. Knechtli, B. Leder, M. Marinkovic, S. Schaefer, R. Sommer and F. Virotta, “The strange quark mass and Lambda parameter of two flavor QCD,” *Nucl. Phys. B* **865**, 397 (2012) [arXiv:1205.5380 [hep-lat]].
- [8] U. Aglietti and Z. Ligeti, “Renormalons and confinement,” *Phys. Lett. B* **364**, 75 (1995) [hep-ph/9503209].
- [9] M. Jezabek, J. H. Kuhn, M. Peter, Y. Sumino and T. Teubner, “The Perturbative QCD potential and the t anti-t threshold,” *Phys. Rev. D* **58**, 014006 (1998) [hep-ph/9802373].
- [10] M. Jezabek, M. Peter and Y. Sumino, “On the relation between QCD potentials in momentum and position space,” *Phys. Lett. B* **428**, 352 (1998) [hep-ph/9803337].
- [11] M. Beneke, “A Quark mass definition adequate for threshold problems,” *Phys. Lett. B* **434**, 115 (1998) [hep-ph/9804241].



- [12] L. S. Brown and W. I. Weisberger, “Remarks on the Static Potential in Quantum Chromodynamics,” *Phys. Rev. D* **20**, 3239 (1979).
- [13] R. Brower, S. Chandrasekharan, J. W. Negele and U. J. Wiese, “QCD at fixed topology,” *Phys. Lett. B* **560**, 64 (2003) [hep-lat/0302005].
- [14] S. Aoki, H. Fukaya, S. Hashimoto and T. Onogi, “Finite volume QCD at fixed topological charge,” *Phys. Rev. D* **76**, 054508 (2007) [arXiv:0707.0396 [hep-lat]].
- [15] M. Lüscher and S. Schaefer, “Lattice QCD without topology barriers,” *JHEP* **1107**, 036 (2011) [arXiv:1105.4749 [hep-lat]].
- [16] W. Bietenholz, I. Hip, S. Shcheredin and J. Volkholz, “A Numerical Study of the 2-Flavour Schwinger Model with Dynamical Overlap Hypercube Fermions,” *Eur. Phys. J. C* **72**, 1938 (2012) [arXiv:1109.2649 [hep-lat]].
- [17] S. Schaefer, “Status and challenges of simulations with dynamical fermions,” *PoS LATTICE* **2012**, 001 (2012) [arXiv:1211.5069 [hep-lat]].
- [18] A. Dromard and M. Wagner, “Extracting hadron masses from fixed topology simulations,” arXiv:1404.0247 [hep-lat].
- [19] A. Billoire, “How heavy must be quarks in order to build Coulombic  $q\bar{q}$  bound states,” *Phys. Lett. B* **92**, 343 (1980).
- [20] A. Pineda, “Determination of the bottom quark mass from the Upsilon(1S) system,” *JHEP* **0106**, 022 (2001) [hep-ph/0105008].
- [21] A. Pineda, “The Static potential: Lattice versus perturbation theory in a renormalon based approach,” *J. Phys. G* **29**, 371 (2003) [hep-ph/0208031].
- [22] A. Laschka, N. Kaiser and W. Weise, “Quark-antiquark potential to order  $1/m$  and heavy quark masses,” *Phys. Rev. D* **83**, 094002 (2011) [arXiv:1102.0945 [hep-ph]].
- [23] F. Karbstein, “Quark-antiquark static energy from a restricted Fourier transform,” *JHEP* **1404**, 144 (2014) [arXiv:1311.7351 [hep-ph]].
- [24] N. Brambilla, A. Vairo, X. Garcia i Tormo and J. Soto, “The QCD static energy at NNNLL,” *Phys. Rev. D* **80**, 034016 (2009) [arXiv:0906.1390 [hep-ph]].
- [25] P. Boucaud *et al.* [ETM Collaboration], “Dynamical twisted mass fermions with light quarks,” *Phys. Lett. B* **650**, 304-311 (2007) [hep-lat/0701012].
- [26] P. Boucaud *et al.* [ETM Collaboration], “Dynamical twisted mass fermions with light quarks: simulation and analysis details,” *Comput. Phys. Commun.* **179**, 695 (2008) [arXiv:0803.0224 [hep-lat]].
- [27] R. Baron *et al.* [ETM Collaboration], “Light meson physics from maximally twisted mass lattice QCD,” *JHEP* **1008**, 097 (2010) [arXiv:0911.5061 [hep-lat]].
- [28] P. Weisz, “Continuum limit improved lattice action for pure Yang-Mills theory. 1,” *Nucl. Phys. B* **212**, 1 (1983).

- [29] R. Frezzotti, P. A. Grassi, S. Sint and P. Weisz [Alpha collaboration], “Lattice QCD with a chirally twisted mass term,” *JHEP* **0108**, 058 (2001) [arXiv:hep-lat/0101001].
- [30] R. Frezzotti and G. C. Rossi, “Chirally improving Wilson fermions. 1.  $\mathcal{O}(a)$  improvement,” *JHEP* **0408**, 007 (2004) [arXiv:hep-lat/0306014].
- [31] R. Frezzotti and G. C. Rossi, “Chirally improving Wilson fermions. 2. Four-quark operators,” *JHEP* **0410**, 070 (2004) [arXiv:hep-lat/0407002].
- [32] A. Shindler, “Twisted mass lattice QCD,” *Phys. Rept.* **461**, 37 (2008) [arXiv:0707.4093 [hep-lat]].
- [33] M. Lüscher, K. Symanzik and P. Weisz, “Anomalies of the free loop wave equation in the WKB approximation,” *Nucl. Phys. B* **173**, 365 (1980).
- [34] M. Lüscher, “Symmetry Breaking Aspects of the Roughening Transition in Gauge Theories,” *Nucl. Phys. B* **180**, 317 (1981).
- [35] M. Donnellan, F. Knechtli, B. Leder and R. Sommer, “Determination of the Static Potential with Dynamical Fermions,” *Nucl. Phys. B* **849**, 45 (2011) [arXiv:1012.3037 [hep-lat]].
- [36] K. Jansen, C. Michael, A. Shindler and M. Wagner [ETM Collaboration], “The static-light meson spectrum from twisted mass lattice QCD,” *JHEP* **0812**, 058 (2008) [arXiv:0810.1843 [hep-lat]].
- [37] A. Sternbeck, “The Infrared behavior of lattice QCD Green’s functions,” hep-lat/0609016.
- [38] R. Sommer, “Scale setting in lattice QCD,” *PoS LATTICE* **2013**, 015 (2014) [arXiv:1401.3270 [hep-lat]].
- [39] W. A. Bardeen, A. J. Buras, D. W. Duke and T. Muta, “Deep Inelastic Scattering Beyond the Leading Order in Asymptotically Free Gauge Theories,” *Phys. Rev. D* **18**, 3998 (1978).
- [40] W. Furmanski and R. Petronzio, “Lepton - Hadron Processes Beyond Leading Order in Quantum Chromodynamics,” *Z. Phys. C* **11**, 293 (1982).
- [41] T. van Ritbergen, J. A. M. Vermaseren and S. A. Larin, “The four-loop beta function in quantum chromodynamics,” *Phys. Lett. B* **400** 379 (1997) [arXiv:hep-ph/9701390].
- [42] T. Appelquist, M. Dine and I. J. Muzinich, “The static limit of quantum chromodynamics,” *Phys. Rev. D* **17** 2074 (1978).
- [43] N. Brambilla, A. Pineda, J. Soto and A. Vairo, “The Infrared behavior of the static potential in perturbative QCD,” *Phys. Rev. D* **60**, 091502 (1999) [hep-ph/9903355].
- [44] F. A. Chishtie and V. Elias, “RG/Pade estimate of the three-loop contribution to the QCD static potential function,” *Phys. Lett. B* **521** 434 (2001) [arXiv:hep-ph/0107052].
- [45] W. Fischler, “Quark-anti-quark potential in QCD,” *Nucl. Phys. B* **129**, 157 (1977).
- [46] M. Peter, “The static quark-anti-quark potential in QCD to three loops,” *Phys. Rev. Lett.* **78**, 602 (1997) [arXiv:hep-ph/9610209].

- [47] M. Peter, “The Static potential in QCD: A Full two loop calculation,” Nucl. Phys. B **501**, 471 (1997) [hep-ph/9702245].
- [48] Y. Schröder, “The static potential in QCD to two loops,” Phys. Lett. B **447**, 321 (1999) [arXiv:hep-ph/9812205].
- [49] A. V. Smirnov, V. A. Smirnov and M. Steinhauser, “Fermionic contributions to the three-loop static potential,” Phys. Lett. B **668**, 293 (2008) [arXiv:0809.1927 [hep-ph]].
- [50] A. V. Smirnov, V. A. Smirnov and M. Steinhauser, “Three-loop static potential,” Phys. Rev. Lett. **104** 112002 (2010) [arXiv:0911.4742 [hep-ph]].
- [51] C. Anzai, Y. Kiyo and Y. Sumino, “Static QCD potential at three-loop order,” Phys. Rev. Lett. **104** 112003 (2010) [arXiv:0911.4335 [hep-ph]].
- [52] A. V. Smirnov, V. A. Smirnov and M. Steinhauser, “Full result for the three-loop static quark potential,” PoS **RADCOR2009**, 075 (2010) [arXiv:1001.2668 [hep-ph]].
- [53] C. Anzai, Y. Kiyo and Y. Sumino, “Violation of Casimir Scaling for Static QCD Potential at Three-loop Order,” Nucl. Phys. B **838**, 28 (2010) [arXiv:1004.1562 [hep-ph]].
- [54] K. G. Chetyrkin, B. A. Kniehl and M. Steinhauser, “Strong coupling constant with flavor thresholds at four loops in the  $\overline{\text{MS}}$  scheme,” Phys. Rev. Lett. **79**, 2184 (1997) [hep-ph/9706430].
- [55] S. Capitani, M. Lüscher, R. Sommer and H. Wittig [ALPHA Collaboration], “Nonperturbative quark mass renormalization in quenched lattice QCD,” Nucl. Phys. B **544**, 669 (1999) [arXiv:hep-lat/9810063].
- [56] J. Beringer *et al.* [Particle Data Group Collaboration], “Review of Particle Physics (RPP),” Phys. Rev. D **86**, 010001 (2012).
- [57] R. Sommer, “A New way to set the energy scale in lattice gauge theories and its applications to the static force and  $\alpha_s$  in SU(2) Yang-Mills theory,” Nucl. Phys. B **411**, 839 (1994) [hep-lat/9310022].
- [58] M. Della Morte, R. Frezzotti, J. Heitger, J. Rolf, R. Sommer and U. Wolff [ALPHA Collaboration], “Computation of the strong coupling in QCD with two dynamical flavors,” Nucl. Phys. B **713**, 378 (2005) [arXiv:hep-lat/0411025].
- [59] A. Sternbeck, E. -M. Ilgenfritz, K. Maltman, M. Müller-Preussker, L. von Smekal, A. G. Williams, “QCD  $\Lambda$  parameter from Landau-gauge gluon and ghost correlations,” PoS **LAT2009**, 210 (2009) [arXiv:1003.1585 [hep-lat]].
- [60] B. Blossier *et al.* [ETM Collaboration], “Ghost-gluon coupling, power corrections and  $\Lambda_{\overline{\text{MS}}}$  from twisted-mass lattice QCD at  $n_f = 2$ ,” Phys. Rev. **D82**, 034510 (2010) [arXiv:1005.5290 [hep-lat]].
- [61] J. -L. Kneur and A. Neveu, “ $\Lambda_{\overline{\text{MS}}}^{\text{QCD}}$  from Renormalization Group Optimized Perturbation,” Phys. Rev. D **85**, 014005 (2012) [arXiv:1108.3501 [hep-ph]].
- [62] K. Cichy, “Quark mass anomalous dimension and  $\Lambda_{\overline{\text{MS}}}$  from the twisted mass Dirac operator spectrum,” arXiv:1311.3572v2 [hep-lat].