

The spectrum of and forces between B mesons from lattice QCD

Seminar Quantentheorie, Friedrich-Schiller-Universität Jena

Marc Wagner

Goethe-Universität Frankfurt am Main, Institut für Theoretische Physik

mwagner@th.physik.uni-frankfurt.de

<http://people.physik.hu-berlin.de/~mcwagner/>

November 17, 2011



Outline

- I will discuss two lattice QCD projects (which are related):
 - (1) **Computation of the spectrum of radially/orbitally excited B mesons.**
 - (2) **Computation of spin/isospin/parity dependent forces between B mesons.**

Part 1

Computation of the spectrum of radially/orbitally excited B mesons

[K. Jansen, C. Michael, A. Shindler and M.W. [ETM Collaboration], JHEP **0812**, 058 (2008)]

[C. Michael, A. Shindler and M.W. [ETM Collaboration], JHEP **1008**, 009 (2010)]


QCD (quantum chromodynamics)

- Quantum field theory of **quarks** (six flavors u, d, s, c, t, b , which differ in **mass**) and **gluons**.
- Part of the standard model explaining the formation of hadrons (mesons = $q\bar{q}$, baryons = $qqq/\bar{q}\bar{q}\bar{q}$) and their masses; essential for decays involving hadrons.
- Definition of QCD by means of an action simple:

$$S = \int d^4x \left(\sum_{f \in \{u,d,s,c,t,b\}} \bar{\psi}^{(f)} \left(\gamma_\mu (\partial_\mu - iA_\mu) + m^{(f)} \right) \psi^{(f)} + \frac{1}{2g^2} \text{Tr} \left(F_{\mu\nu} F_{\mu\nu} \right) \right)$$

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - i[A_\mu, A_\nu].$$

- However, no analytical solutions for low energy QCD observables, e.g. hadron masses, known, because of the absence of any small parameter (i.e. perturbation theory not applicable).



B mesons, static-light mesons


- *B* meson: a meson made from a heavy *b* quark ($m_b \approx 4200$ MeV) and a light *u*, *d* or *s* quark ($m_l \lesssim 100$ MeV), e.g. $B = \{\bar{b}u, \bar{b}d\}$, $B_s = \bar{b}s$.
- Static limit, i.e. $m_b \rightarrow \infty$:

- No interactions involving the static quark spin.
- Classify states according to parity \mathcal{P} and half-integer total angular momentum of the light cloud j .

- m_b finite, but heavy:

- Classify states according to parity \mathcal{P} and total angular momentum J .

$j^{\mathcal{P}}$ (static-light)	$J^{\mathcal{P}}$ (finite m_b)
$(1/2)^- \equiv S$	$0^- \equiv B_{(s)}$ $1^- \equiv B_{(s)}^*$
$(1/2)^+ \equiv P_-$	$0^+ \equiv B_{(s)0}^*$ (not in PDG) $1^+ \equiv B_{(s)1}^*$ (not in PDG)
$(3/2)^+ \equiv P_+$	$1^+ \equiv B_{(s)1}$ $2^+ \equiv B_{(s)2}^*$
$(3/2)^- \equiv D_-$	1^- (no experiment) 2^- (no experiment)
$(5/2)^- \equiv D_+$	2^- (no experiment) 3^- (no experiment)
$(5/2)^+ \equiv F_-$	2^+ (no experiment) 3^+ (no experiment)
...	...



How to compute M (static-light meson)? (1)

- Let $\mathcal{O}(\mathbf{x})$ be a suitable “static-light meson creation operator”, i.e. an operator such that $\mathcal{O}(\mathbf{x})|\Omega\rangle$ is a state containing a static-light meson at position \mathbf{x} ($|\Omega\rangle$: vacuum).
- More precisely: ... an operator such that $\mathcal{O}(\mathbf{x})|\Omega\rangle$ has the same quantum numbers (j^P , flavor) as the static-light meson of interest.
- Determine the mass of the ground state of the corresponding static-light meson from the exponential behavior of the corresponding correlation function \mathcal{C} at large Euclidean times T :

$$\begin{aligned}\mathcal{C}(T) &= \langle \Omega | (\mathcal{O}(\mathbf{x}, T))^\dagger \mathcal{O}(\mathbf{x}, 0) | \Omega \rangle = \langle \Omega | e^{+HT} (\mathcal{O}(\mathbf{x}, 0))^\dagger e^{-HT} \mathcal{O}(\mathbf{x}, 0) | \Omega \rangle = \\ &= \sum_n \left| \langle n | \mathcal{O}(\mathbf{x}, 0) | \Omega \rangle \right|^2 \exp \left(- (E_n - E_\Omega) T \right) \approx \quad (\text{for } T \gg 1) \\ &\approx \left| \langle 0 | \mathcal{O}(\mathbf{x}, 0) | \Omega \rangle \right|^2 \exp \left(- \underbrace{(E_0 - E_\Omega)}_{M(\text{static-light meson})} T \right).\end{aligned}$$

How to compute M (static-light ...)

- General form of a static-light meson creation operator:


$$\mathcal{O}(\mathbf{x}) = \bar{Q}(\mathbf{x}) \int d\hat{\mathbf{n}} \Gamma(\hat{\mathbf{n}}) U(\mathbf{x}; \mathbf{x} + d\hat{\mathbf{n}}) q(\mathbf{x} + d\hat{\mathbf{n}}).$$

- $\bar{Q}(\mathbf{x})$ creates an infinitely heavy i.e. static antiquark at position \mathbf{x} .
- $q(\mathbf{x} + d\hat{\mathbf{n}})$ creates a light quark at position $\mathbf{x} + d\hat{\mathbf{n}}$ separated by a distance d from the static antiquark.
- The spatial parallel transporter

$$U(\mathbf{x}; \mathbf{x} + d\hat{\mathbf{n}}) = P \left\{ \exp \left(+i \int_{\mathbf{x}}^{\mathbf{x}+d\hat{\mathbf{n}}} dz_j A_j(\mathbf{z}) \right) \right\}$$

connects the antiquark and the quark in a gauge invariant way via gluons.


- The integration over the unit sphere $\int d\hat{\mathbf{n}}$ combined with a suitable weight factor $\Gamma(\hat{\mathbf{n}})$ yields well defined total angular momentum J and parity \mathcal{P} ($\Gamma(\hat{\mathbf{n}})$ is a combination of spherical harmonics [\rightarrow angular momentum] and γ -matrices [\rightarrow spin]; Wigner-Eckart theorem).



How to compute M (static-light ...)

- General form of a static-light meson creation operator:

$$\mathcal{O}(\mathbf{x}) = \bar{Q}(\mathbf{x}) \int d\hat{\mathbf{n}} \Gamma(\hat{\mathbf{n}}) U(\mathbf{x}; \mathbf{x} + d\hat{\mathbf{n}}) q(\mathbf{x} + d\hat{\mathbf{n}}).$$



- List of operators (J : total angular momentum; j : total angular momentum of the light cloud; \mathcal{P} : parity):

$\Gamma(\hat{\mathbf{n}})$	$J^{\mathcal{P}}$	$j^{\mathcal{P}}$	O_h	lattice $j^{\mathcal{P}}$	notation
$\gamma_5, \gamma_5 \gamma_j \hat{n}_j$ $1, \gamma_j \hat{n}_j$	$0^- [1^-]$ $0^+ [1^+]$	$(1/2)^-$ $(1/2)^+$	A_1	$(1/2)^-, (7/2)^-, \dots$ $(1/2)^+, (7/2)^+, \dots$	S P_-
$\gamma_1 \hat{n}_1 - \gamma_2 \hat{n}_2$ (and cyclic) $\gamma_5 (\gamma_1 \hat{n}_1 - \gamma_2 \hat{n}_2)$ (and cyclic)	$2^+ [1^+]$ $2^- [1^-]$	$(3/2)^+$ $(3/2)^-$	E	$(3/2)^+, (5/2)^+, \dots$ $(3/2)^-, (5/2)^-, \dots$	P_+ D_{\pm}
$\gamma_1 \hat{n}_2 \hat{n}_3 + \gamma_2 \hat{n}_3 \hat{n}_1 + \gamma_3 \hat{n}_1 \hat{n}_2$ $\gamma_5 (\gamma_1 \hat{n}_2 \hat{n}_3 + \gamma_2 \hat{n}_3 \hat{n}_1 + \gamma_3 \hat{n}_1 \hat{n}_2)$	$3^- [2^-]$ $3^+ [2^+]$	$(5/2)^-$ $(5/2)^+$	A_2	$(5/2)^-, (7/2)^-, \dots$ $(5/2)^+, (7/2)^+, \dots$	D_+ F_{\pm}

Lattice QCD (1)


- Goal: compute correlation functions $\mathcal{C}(T)$ of the previously discussed static-light meson creation operators (the corresponding meson masses can directly be read off from their exponential decays).
- Use the path integral formulation of QCD,

$$\begin{aligned}\mathcal{C}(T) &= \langle \Omega | \left(\mathcal{O}(\mathbf{x}, T) \right)^\dagger \mathcal{O}(\mathbf{x}, 0) | \Omega \rangle = \\ &= \frac{1}{Z} \int \left(\prod_f D\psi^{(f)} D\bar{\psi}^{(f)} \right) DA_\mu \left(\mathcal{O}(\mathbf{x}, T) \right)^\dagger \mathcal{O}(\mathbf{x}, 0) e^{-S[\psi^{(f)}, \bar{\psi}^{(f)}, A_\mu]}.\end{aligned}$$

- $|\Omega\rangle$: ground state/vacuum.
- $(\mathcal{O}(\mathbf{x}, T))^\dagger \mathcal{O}(\mathbf{x}, 0)$: function of the quark and gluon fields (cf. previous slides).
- $\int (\prod_f D\psi^{(f)} D\bar{\psi}^{(f)}) DA_\mu$: integral over all possible quark and gluon field configurations $\psi^{(f)}(\mathbf{x}, t)$ and $A_\mu(\mathbf{x}, t)$.
- $e^{-S[x]}$: weight factor containing the QCD action.

Lattice QCD (2)

- Numerical implementation of the path integral formalism in QCD:
 - Discretize spacetime with sufficiently small lattice spacing
 $a \approx 0.05 \text{ fm} \dots 0.10 \text{ fm}$
→ “continuum physics”.
 - “Make spacetime periodic” with sufficiently large extension
 $L \approx 2.0 \text{ fm} \dots 4.0 \text{ fm}$ (4-dimensional torus)
→ “no finite size effects”.



Lattice QCD (3)

- Numerical implementation of the path integral formalism in QCD:
 - After discretization the path integral becomes an ordinary multidimensional integral:


$$\int D\psi D\bar{\psi} DA \dots \rightarrow \prod_{x_\mu} \left(\int d\psi(x_\mu) d\bar{\psi}(x_\mu) dU(x_\mu) \right) \dots$$

- Typical present-day dimensionality of a discretized QCD path integral:
 - * x_μ : $32^4 \approx 10^6$ lattice sites.
 - * $\psi = \psi_A^{a,(f)}$: 24 quark degrees of freedom for every flavor ($\times 2$ particle/antiparticle, $\times 3$ color, $\times 4$ spin), 2 flavors.
 - * $U = U_\mu^{ab}$: 32 gluon degrees of freedom ($\times 8$ color, $\times 4$ spin).
 - * In total: $32^4 \times (2 \times 24 + 32) \approx \mathbf{83} \times \mathbf{10^6}$ dimensional integral.
- standard approaches for numerical integration not applicable
- sophisticated algorithms mandatory (stochastic integration techniques, so-called Monte-Carlo algorithms).







Lattice setup

- Various lattice spacings: $a \approx 0.051 \text{ fm}$, 0.064 fm , 0.080 fm (corresponding to $48^3 \times 96$, $32^3 \times 64$, $24^3 \times 48$ lattice sites).
- Lattice extensions: $L \approx 2.45 \text{ fm}$, 2.05 fm , 1.92 fm (periodic boundary conditions).
- Many different pion mass: $m_{\text{PS}} \approx 284 \dots 637 \text{ MeV}$.


Masses of B and B_s mesons (1)



- Compute static-light meson masses (B/B_s mesons with $m_b \rightarrow \infty$) for different light u/d quark masses and different lattice spacings:
 - Different u/d quark masses to extrapolate to the physical u/d quark mass (due to technical reasons $m_{PS}^{(\text{lattice})} \gtrsim 284$ MeV, $m_{PS}^{\text{physical}} \approx 135$ MeV).
 - Different lattice spacings to extrapolate to the continuum.
 - Horizontal axis: pion mass ($m_{PS}^{(\text{lattice})}$)².
 - Vertical axis: $M(j^P) - M((1/2)^-)$ mass difference between radially and orbitally excited “ B mesons” (B_0^* , B_1^* , B_1 , B_2^* , ...) and the “ground state B meson” ($B/B^* \equiv j^P = (1/2)^-$) ... analogous for “ B_s mesons”.



Masses of B and B_s mesons (2)




- Summary of the computed static-light meson spectrum:

j^P	alternative notation	B mesons ($\bar{b}u$ or $\bar{b}d$): $M(j^P) - M((1/2)^-)$ in MeV	B_s mesons ($\bar{b}s$): $M(j^P) - M((1/2)^-)$ in MeV
$(1/2)^+$	P_-	406(19)	413(12)
$(3/2)^+$	P_+	516(18)	504(12)
$(3/2)^-, (5/2)^-$	D_\pm	870(27)	770(26)
$(5/2)^-$	D_+	930(28)	960(24)
$(5/2)^+, (7/2)^+$	F_\pm	1196(30)	1179(37)
$(1/2)^-$	S^*	755(16)	751(26)

- Motivation/achievements:
 - Continuum limit (among the first).
 - Dependence on the light u/d sea quark mass (for the first time).
 - Valuable input for model builders (e.g. no reversal of $M(P_-)$ and $M(P_+)$, ...).

Masses of B and B_s mesons (3)



- Comparison to experimental results:
 - Extrapolation to the physical (finite) b quark mass $m_B \approx 4200$ MeV:
 - * Use rather precise experimental results for c quarks, i.e. D mesons.
 - * Assume that Heavy Quark Effective Theory (HQET) up to $\mathcal{O}(1/m_Q)$ is “valid” down to the physical charm quark mass.
 - * Amounts to “reincluding” hyperfine splitting.

	$M - M(B)$ in MeV			$M - M(B_s)$ in MeV	
name	lattice	experiment	name	lattice	experiment
B_0^*	443(21)		B_{s0}^*	391(8)	
B_1^*	460(22)		B_{s1}^*	440(8)	
B_1	530(12)	444(2)	B_{s1}	526(8)	463(1)
B_2^*	543(12)	464(5)	B_{s2}^*	539(8)	473(1)
B_J^*		418(8)	B_{sJ}^*		487(15)

- Difference between lattice and experimental results: scale setting problem?

Part 2


Computation of spin/isospin/parity dependent forces between B mesons.

[M. W. [ETM Collaboration], PoS **LATTICE2010**, 162 (2010)]

[M. W. [ETM Collaboration], Acta Phys. Pol. B Proceedings Supplement, Vol. 4, No. 4, 2011, page 747]

Introduction (1)

- Goal: compute the potential of (or equivalently the force between) two B mesons from first principles by means of lattice QCD:
 - Treat the b quark in the static approximation.
 - Consider only pseudoscalar/vector mesons ($j^{\mathcal{P}} = (1/2)^-$, denoted by S , PDG: B, B^*) and scalar/pseudovector mesons ($j^{\mathcal{P}} = (1/2)^+$, denoted by P_- , PDG: B_0^*, B_1^*), which are among the lightest static-light mesons.
 - Study the dependence of the mesonic potential $V(R)$ on
 - * the light quark flavor u and/or d (isospin),
 - * the light quark spin (the static quark spin is irrelevant),
 - * the type of the meson S and/or P_- .



Introduction (2)


- Motivation:

- First principles computation of a hadronic force.
- Possible application: determine, whether two B mesons may form bound states (tetraquarks).
- Until now
 - * it has mainly been studied in the quenched approximation,
 - * only pseudoscalar (S), but no scalar (P_-) B mesons have been considered.

[C. Michael and P. Pennanen [UKQCD Collaboration], Phys. Rev. D **60**, 054012 (1999)]

[W. Detmold, K. Orginos and M. J. Savage, Phys. Rev. D **76**, 114503 (2007)]

[G. Bali and M. Hetzenegger, PoS LATTICE2010, 142 (2010)]




(Pseudo)scalar B mesons

- Symmetries and quantum numbers of static-light mesons:
 - Isospin: $I = 1/2$, $I_z = \pm 1/2$, i.e. $B \equiv \bar{Q}u$ or $B \equiv \bar{Q}d$.
 - Parity: $\mathcal{P} = \pm$,
 - * $\mathcal{P} = - \equiv S$ (wave),
 - * $\mathcal{P} = + \equiv P_-$ (wave).
 - Rotations:
 - * Light cloud angular momentum $j = 1/2$ (for S and P_-), $j_z = \pm 1/2$.
 - * Static quark spin: irrelevant (static quarks can also be treated as spinless color charges).
- Examples of static-light meson creation operators:
 - $\bar{Q}\gamma_5 q$ (pseudoscalar, i.e. S), $q \in \{u, d\}$,
 - $\bar{Q}q$ (scalar, i.e. P_-)

(j_z is not well-defined, when using these operators).

BB systems (1)

- Symmetries and quantum numbers of a pair of static-light mesons (separated along the z -axis):
 - Isospin: $I = 0, 1, I_z = -1, 0, +1$.
 - Rotations around the z -axis:
 - * Angular momentum of the light degrees of freedom $j_z = -1, 0, +1$.
 - * Static quark spin: irrelevant (static quarks can also be treated as spinless color charges).
 - Parity: $\mathcal{P} = \pm$.
 - If $j_z = 0$, reflection along the x -axis: $\mathcal{P}_x = \pm$.
 - Instead of using $j_z = \pm 1$ one can also label states by $|j_z| = 1, \mathcal{P}_x = \pm$.
- Label BB states by $(I, I_z, |j_z|, \mathcal{P}, \mathcal{P}_x)$.




BB systems (2)

- To extract the potential(s) of a given sector (characterized by $(I, I_z, |j_z|, \mathcal{P}, \mathcal{P}_x)$), compute the temporal correlation function of the trial state

$$(\mathcal{C}\Gamma)_{AB} \left(\bar{Q}_C(-R/2) q_A^{(1)}(-R/2) \right) \left(\bar{Q}_C(+R/2) q_B^{(2)}(+R/2) \right) |\Omega\rangle,$$

where

- $\mathcal{C} = \gamma_0 \gamma_2$ (charge conjugation matrix),
- $q^{(1)} q^{(2)} \in \{ud - du, uu, dd, ud + du\}$ (isospin I, I_z),
- Γ is an arbitrary combination of γ matrices (spin $|j_z|$, parity $\mathcal{P}, \mathcal{P}_x$).



BB systems (3)

- *BB* creation operators for $I_z = +1$: 16 operators of type

$$(\mathcal{C}\Gamma)_{AB} \left(\bar{Q}_C(-R/2) q_A^{(u)}(-R/2) \right) \left(\bar{Q}_C(+R/2) q_B^{(u)}(+R/2) \right).$$

Γ	$ j_z , \mathcal{P}, \mathcal{P}_x$
1	0, -, -
$\gamma_0 \gamma_5$	0, +, +
γ_5	0, +, +
γ_0	0, +, -
γ_3	0, -, -
$\gamma_0 \gamma_3 \gamma_5$	0, +, +
$\gamma_3 \gamma_5$	0, -, +
$\gamma_0 \gamma_3$	0, -, -
γ_1	1, -, +
$\gamma_0 \gamma_1 \gamma_5$	1, +, -
$\gamma_1 \gamma_5$	1, -, -
$\gamma_0 \gamma_1$	1, -, +
...	...

BB systems (4)

- *BB* creation operators for $I_z = 0$: 32 operators of type

$$(\mathcal{C}\Gamma)_{AB} \left(\bar{Q}_C(-R/2) q_A^{(u)}(-R/2) \right) \left(\bar{Q}_C(+R/2) q_B^{(d)}(+R/2) \right) \pm (u \leftrightarrow d).$$


Γ, \pm	$ j_z , I, \mathcal{P}, \mathcal{P}$
$\gamma_5, -$	0, 0, -, +
$\gamma_0, -$	0, 0, -, -
1, -	0, 0, +, -
$\gamma_0\gamma_5, -$	0, 0, -, +
$\gamma_3\gamma_5, -$	0, 0, +, +
$\gamma_0\gamma_3, -$	0, 0, +, -
$\gamma_3, -$	0, 0, +, -
$\gamma_0\gamma_3\gamma_5, -$	0, 0, -, +
$\gamma_5, +$	0, 1, +, +
$\gamma_0, +$	0, 1, +, -
1, +	0, 1, -, -
$\gamma_0\gamma_5, +$	0, 1, +, +
...	...

Lattice setup

- Lattice spacing: $a \approx 0.079$ fm.
- Lattice extension: $L \approx 1.90$ fm (periodic boundary conditions).
- Pion mass: $m_{\text{PS}} \approx 340$ MeV.


Discussion of results (1)

- Four “types of potentials”:
 - Two attractive, two repulsive.
 - Two have asymptotic values, which are larger by ≈ 400 MeV.
- There are cases, where two potentials with identical quantum numbers are completely different (i.e. of different type)
 - at least one of the corresponding trial states must have very small ground state overlap
 - physical understanding, i.e. interpretation of trial states needed.



Discussion of results (2)

- Expectation at large meson separation R : $V(R) \approx 2 \times \text{meson mass}$.
 - Potentials with smaller asymptotic value at $\approx 2 \times m(S)$.
 - $m(P_-) - m(S) \approx 400 \text{ MeV}$: approximately the observed difference between different types of potentials.
- Two types correspond to $S \leftrightarrow S$ potentials.
- Two types correspond to $S \leftrightarrow P_-$ potentials.
- Can this be understood in detail on the level of the used BB creation operators?



Discussion of results (3)

- Express the BB creation operators in terms of static-light meson creation operators (use suitable spin and parity projectors for the light quarks).

– Examples:

$$* uu, \Gamma = 1 \quad \rightarrow \quad \mathcal{P} = -, \mathcal{P}_x = -:$$


$$(\mathcal{C}1)_{AB} \left(\bar{Q}_C(-R/2) q_A^{(u)}(-R/2) \right) \left(\bar{Q}_C(+R/2) q_B^{(u)}(+R/2) \right) \propto \\ \propto S_\uparrow P_\downarrow - S_\downarrow P_\uparrow + P_\uparrow S_\downarrow - P_\downarrow S_\uparrow.$$

$$* uu, \Gamma = \gamma_3 \quad \rightarrow \quad \mathcal{P} = -, \mathcal{P}_x = -:$$

$$(\mathcal{C}\gamma_3)_{AB} \left(\bar{Q}_C(-R/2) q_A^{(u)}(-R/2) \right) \left(\bar{Q}_C(+R/2) q_B^{(u)}(+R/2) \right) \propto \\ \propto S_\uparrow S_\downarrow + S_\downarrow S_\uparrow - P_\uparrow P_\downarrow - P_\downarrow P_\uparrow.$$


– SS/SP_- content and asymptotic values in agreement for all 64 correlation functions/
potentials

→ asymptotic differences understood.



Discussion of results (4)

- Is there a general rule, about when a potential is repulsive and when attractive?



– $S \leftrightarrow S$ potentials:

- * $(I = 0, s = 0)$ or $(I = 1, s = 1)$, i.e. $I = s \rightarrow$ attractive
 - * $(I = 0, s = 1)$ or $(I = 1, s = 0)$, i.e. $I \neq s \rightarrow$ repulsive
- (s : combined angular momentum of the two mesons).

- * **Example:** $uu, \Gamma = \gamma_3 \rightarrow \mathcal{P} = -, \mathcal{P}_x = -:$

$$\begin{aligned}
 & (\mathcal{C}\gamma_3)_{AB} \left(\bar{Q}_C(-R/2) q_A^{(u)}(-R/2) \right) \left(\bar{Q}_C(+R/2) q_B^{(u)}(+R/2) \right) \propto \\
 & \propto S_\uparrow S_\downarrow + S_\downarrow S_\uparrow - P_\uparrow P_\downarrow - P_\downarrow P_\uparrow.
 \end{aligned}$$

i.e. $I = 1, s = 1$; the numerically obtained potential is attractive, i.e. in agreement with the above stated rule.

- * All 32 $S \leftrightarrow S$ correlation functions/potentials fulfill the rule.
- * Agreement with similar quenched lattice studies.

[C. Michael and P. Pennanen [UKQCD Collaboration], Phys. Rev. D **60**, 054012 (1999)]

[W. Detmold, K. Orginos and M. J. Savage, Phys. Rev. D **76**, 114503 (2007)]

Discussion of results (5)

– $S \leftrightarrow P_-$ potentials:

- * Do not obey the above stated rule.


- * It can, however, easily be generalized by including parity, i.e. symmetry or antisymmetry under exchange of S and P_- :
 - trial state symmetric under meson exchange \rightarrow attractive
 - trial state antisymmetric under meson exchange \rightarrow repulsive (meson exchange \equiv exchange of flavor, spin and parity).

- * **Example:** $uu, \Gamma = \gamma_0 \rightarrow \mathcal{P} = +, \mathcal{P}_x = -$:

$$\begin{aligned}
 (\mathcal{C}\gamma_0)_{AB} \left(\bar{Q}_C(-R/2) q_A^{(u)}(-R/2) \right) \left(\bar{Q}_C(+R/2) q_B^{(u)}(+R/2) \right) &\propto \\
 \propto S_\uparrow P_\downarrow - S_\downarrow P_\uparrow - P_\uparrow S_\downarrow + P_\downarrow S_\uparrow, &
 \end{aligned}$$


i.e. $I = 1$ (symmetric), $s = 0$ (antisymmetric), antisymmetric with respect to $S \leftrightarrow P_-$; the numerically obtained potential is attractive, i.e. in agreement with the above stated general rule.

- * All 32 $S \leftrightarrow P_-$ correlation functions/potentials (and all 32 $S \leftrightarrow S$ correlation functions/potentials) fulfill the generalized rule.



Discussion of results (6)

- Improvements after having understood the extraction and interpretation of BB potentials from single correlation functions:
 - Linearly combine BB operators to either eliminate $P_- \leftrightarrow P_-$ or $S \leftrightarrow S$ combinations.
 - Example:
 - $ud - du, \Gamma = \gamma_5 \rightarrow -S_\uparrow S_\downarrow + S_\downarrow S_\uparrow - P_\uparrow P_\downarrow + P_\downarrow P_\uparrow$
 - $ud - du, \Gamma = \gamma_0 \gamma_5 \rightarrow -S_\uparrow S_\downarrow + S_\downarrow S_\uparrow + P_\uparrow P_\downarrow - P_\downarrow P_\uparrow$
 - use $\gamma_5 + \gamma_0 \gamma_5$ to obtain a better signal for the $S \leftrightarrow S$ potential
 - use $\gamma_5 - \gamma_0 \gamma_5$ to extract the $P_- \leftrightarrow P_-$ potential.



Discussion of results (7)

- Improvements after having understood the extraction and interpretation of BB potentials from single correlation functions:
 - Use correlation matrices instead of single correlation functions to avoid mixing with BB states of lower energy, which is present, because
 - * although the product of two specific B meson creation operators closely resembles the corresponding BB state, it will still have a non-vanishing overlap to BB states corresponding to B mesons with different isospin, spin and/or parity,
 - * twisted mass lattice QCD explicitly breaks isospin and parity (the breaking is proportional to the lattice spacing a ; isospin and parity will be restored in the continuum limit).

Summary of BB states and degeneracies

- Two B mesons, each can have $I_z = \pm 1/2$, $j_z = \pm 1/2$, $\mathcal{P} = \pm$
 $\rightarrow 8 \times 8 = 64$ states.

- $S \leftrightarrow S$ potentials:

– Attractive: $\underbrace{1}_{I=0,|j_z|=0} \oplus \underbrace{3}_{I=1,|j_z|=0} \oplus \underbrace{6}_{I=1,|j_z|=1}$ (10 states).

– Repulsive: $\underbrace{1}_{I=0,|j_z|=0} \oplus \underbrace{3}_{I=1,|j_z|=0} \oplus \underbrace{2}_{I=0,|j_z|=1}$ (6 states).

- $S \leftrightarrow P_-$ potentials:

– Attractive: $\underbrace{1 \oplus 1 \oplus 3 \oplus 3}_{|j_z|=0} \oplus \underbrace{2 \oplus 6}_{|j_z|=1}$ (16 states).

– Repulsive: $\underbrace{1 \oplus 1 \oplus 3 \oplus 3}_{|j_z|=0} \oplus \underbrace{2 \oplus 6}_{|j_z|=1}$ (16 states).




- $P_- \leftrightarrow P_-$ potentials: identical to $S \leftrightarrow S$ potentials.

- In total 24 different potentials.

Attractive $S \leftrightarrow S$ potentials

- Attractive $S \leftrightarrow S$ potentials are relevant, when trying to determine, whether BB may form a bound state.

- Three different attractive $S \leftrightarrow S$ potentials: $\underbrace{1}_{I=0, |j_z|=0} \oplus \underbrace{3}_{I=1, |j_z|=0} \oplus \underbrace{6}_{I=1, |j_z|=1}$.



Summary, conclusions, future plans (1)

- Computation of BB potentials (arbitrary flavor, spin and parity) with “light” dynamical quarks ($m_{\text{PS}} \approx 340 \text{ MeV}$).
 - Qualitative agreement with existing quenched results for $S \leftrightarrow S$ potentials.
 - First lattice computation of $S \leftrightarrow P_-$ and $P_- \leftrightarrow P_-$ potentials.
 - Clear statements about whether a potential of a given channel is attractive or repulsive.
- Statistical accuracy problematic (exponentially decaying correlation functions are quickly lost in statistical noise):
 - Reasonable accuracy for attractive $S \leftrightarrow S$ potentials (interesting, when trying to determine, whether BB may form a bound state).
 - Other (higher) potentials:
 - BB potentials are extracted at rather small temporal separations
 - slight contamination from excited states cannot be excluded.

Summary, conclusions, future plans (2)

- Further plans and possibilities:
 - Other values of the lattice spacing, the spacetime volume and/or the u/d quark mass.
 - Partially quenched computations, to obtain $B_s B_s$ and/or $B_s B$ potentials.
 - Improve lattice meson potentials at small separations (where the suppression of UV fluctuations due to the lattice cutoff yields wrong results) with corresponding perturbative potentials.
 - Use lattice meson potentials to study, whether BB may form a bound state.