

# Comparing meson-meson and diquark-antidiquark creation operators for a $\bar{b}\bar{b}ud$ tetraquark

“38th International Symposium on Lattice Field Theory”

Marc Wagner

Goethe-Universität Frankfurt, Institut für Theoretische Physik

[mwagner@itp.uni-frankfurt.de](mailto:mwagner@itp.uni-frankfurt.de)

<https://itp.uni-frankfurt.de/~mwagner/>

in collaboration with Pedro Bicudo, Antje Peters, Sebastian Velten

July 28, 2021




[P. Bicudo, A. Peters, S. Velten, M.W., Phys. Rev. D **103**, 114506 (2021) [arXiv:2101.00723]]

# Basic idea: lattice QCD + BO


- Study heavy-heavy-light-light tetraquarks  $\bar{b}\bar{b}qq$  in two steps.
  - (1) Compute potentials of two static quarks  $\bar{b}\bar{b}$  in the presence of two lighter quarks  $qq$  ( $q \in \{u, d, s, c\}$ ) using lattice QCD.
  - (2) Check, whether these potentials are sufficiently attractive to host bound states or resonances ( $\rightarrow$  tetraquarks) by using techniques from quantum mechanics and scattering theory.

((1) + (2)  $\rightarrow$  Born-Oppenheimer approximation).

step 1



step 2



$\rightarrow$  existence of a tetraquark ... or not

# Previous work on $b\bar{b}qq$ tetraquarks


- Lattice QCD static potentials and Born-Oppenheimer approximation.  
[W. Detmold, K. Orginos, M. J. Savage, Phys. Rev. D **76**, 114503 (2007) [arXiv:hep-lat/0703009]]  
[M.W., PoS **LATTICE2010**, 162 (2010) [arXiv:1008.1538]]  
[G. Bali, M. Hetzenegger, PoS **LATTICE2010**, 142 (2010) [arXiv:1011.0571]]  
[P. Bicudo, M.W., Phys. Rev. D **87**, 114511 (2013) [arXiv:1209.6274]]  
[Z. S. Brown, K. Orginos, Phys. Rev. D **86**, 114506 (2012) [arXiv:1210.1953]]  
[P. Bicudo, K. Cichy, A. Peters, B. Wagenbach, M.W., Phys. Rev. D **92**, 014507 (2015) [arXiv:1505.00613]]  
[P. Bicudo, K. Cichy, A. Peters, M.W., Phys. Rev. D **93**, 034501 (2016) [arXiv:1510.03441]]  
[P. Bicudo, J. Scheunert, M.W., Phys. Rev. D **95**, 034502 (2017) [arXiv:1612.02758]]  
[P. Bicudo, M. Cardoso, A. Peters, M. Pflaumer, M.W., Phys. Rev. D **96**, 054510 (2017) [arXiv:1704.02383]]  
[P. Bicudo, A. Peters, S. Velten, M.W., Phys. Rev. D **103**, 114506 (2021) [arXiv:2101.00723]]
- Full lattice QCD ( $b$  quarks with Non Relativistic QCD) [list not complete]:  
[A. Francis, R. J. Hudspith, R. Lewis, K. Maltman, Phys. Rev. Lett. **118**, 142001 (2017) [arXiv:1607.05214 [hep-lat]]]  
[P. Junnarkar, N. Mathur, M. Padmanath, Phys. Rev. D **99**, 034507 (2019) [arXiv:1810.12285 [hep-lat]]]  
[L. Leskovec, S. Meinel, M. Pflaumer, M.W., Phys. Rev. D **100**, 014503 (2019) [arXiv:1904.04197 [hep-lat]]]
- Other approaches: quark models, effective field theories, QCD sum rules ... [list not complete]:  
[M. Karliner, J. L. Rosner, Phys. Rev. Lett. **119**, 202001 (2017) [arXiv:1707.07666]]  
[E. J. Eichten, C. Quigg, Phys. Rev. Lett. **119**, 202002 (2017) [arXiv:1707.09575]]  
[Z. G. Wang, Acta Phys. Polon. B **49**, 1781 (2018) [arXiv:1708.04545]]  
[W. Park, S. Noh, S. H. Lee, Acta Phys. Polon. B **50**, 1151-1157 (2019) [arXiv:1809.05257]]  
[B. Wang, Z. W. Liu, X. Liu, Phys. Rev. D **99**, 036007 (2019) [arXiv:1812.04457]]  
[M. Z. Liu, T. W. Wu, M. Pavon Valderrama, J. J. Xie, L. S. Geng, Phys. Rev. D **99**, 094018 (2019) [arXiv:1902.03044]]

# Outline

- $\bar{b}bqq$  /  $BB$  potentials.
- Stable  $\bar{b}bqq$  tetraquarks.
- Structure of a  $\bar{b}bqq$  tetraquark with quantum numbers  $I(J^P) = 0(1^+)$  (meson-meson versus diquark-antidiquark structure).

# $\bar{b}\bar{b}qq$ / $BB$ potentials (1)

- At large  $\bar{b}\bar{b}$  separation  $r$ , the four quarks will form two static-light mesons  $\bar{b}q$  and  $\bar{b}q$ .
  - Spins of static antiquarks  $\bar{b}\bar{b}$  are irrelevant (they do not appear in the Hamiltonian).
  - Compute and study the dependence of  $\bar{b}\bar{b}$  potentials in the presence of  $qq$  on
    - the “light” quark flavors  $q \in \{u, d, s, c\}$  (isospin, flavor),
    - the “light” quark spin (the static quark spin is irrelevant),
    - the type of the meson  $B, B^*$  and/or  $B_0^*, B_1^*$  (parity).
- Many different channels: attractive as well as repulsive, different asymptotic values ...



# $\bar{b}\bar{b}qq$ / $BB$ potentials (2)

- To determine potentials, compute temporal correlation functions of operators

$$\mathcal{O}_{BB,\Gamma} = 2N_{BB}(\mathcal{C}\Gamma)_{AB}(\mathcal{C}\tilde{\Gamma})_{CD} \left( \bar{Q}_C^a(-\mathbf{r}/2)\psi_A^{(f)a}(-\mathbf{r}/2) \right) \left( \bar{Q}_D^b(+\mathbf{r}/2)\psi_B^{(f')b}(+\mathbf{r}/2) \right).$$

- The most attractive potential of a  $B^{(*)}B^*$  meson pair has  $(I, |j_z|, P, P_x) = (0, 0, +, -)$ :


- $C = \gamma_0\gamma_2$  (charge conjugation matrix).
- $\psi^{(f)}\psi^{(f')} = ud - du$ ,  $\Gamma \in \{(1 + \gamma_0)\gamma_5, (1 - \gamma_0)\gamma_5\}$ .
- $\bar{Q}\bar{Q} = \bar{b}\bar{b}$ ,  $\tilde{\Gamma} \in \{(1 + \gamma_0)\gamma_5, (1 + \gamma_0)\gamma_j\}$  (irrelevant).

- Parameterize lattice results by

$$V_{qq,j_z,\mathcal{P},\mathcal{P}_x}(r) = -\frac{\alpha}{r} \exp\left(-\left(\frac{r}{d}\right)^p\right) + V_0$$

(1-gluon exchange at small  $r$ ; color screening at large  $r$ ).

[P. Bicudo, K. Cichy, A. Peters, M.W., Phys. Rev. D **93**, 034501 (2016) [arXiv:1510.03441]]




# Stable $\bar{b}\bar{b}qq$ tetraquarks

- Solve the Schrödinger equation for the relative coordinate of the heavy quarks  $\bar{b}\bar{b}$  using the previously computed  $\bar{b}\bar{b}qq / BB$  potentials,

$$\left( \frac{1}{m_b} \left( -\frac{d^2}{dr^2} + \frac{L(L+1)}{r^2} \right) + V_{qq,j_z, \mathcal{P}, \mathcal{P}_x}(r) - 2m_{sl} \right) R(r) = ER(r).$$

- Possibly existing bound states, i.e.  $E < 0$ , indicate stable  $\bar{b}\bar{b}qq$  tetraquarks.
- There is a bound state for orbital angular momentum  $L = 0$  of  $\bar{b}\bar{b}$ :
  - Binding energy  $-E = 38(18)$  MeV with respect to the  $BB^*$  threshold.
  - Quantum numbers:  $I(J^P) = 0(1^+)$ .
- No further bound states.

[P. Bicudo, M.W., Phys. Rev. D 87, 114511 (2013) [arXiv:1209.6274]]



# Structure of the $\bar{b}\bar{b}qq$ tetraquark (1)

- Two types of operators, which probe the same sector:  
[P. Bicudo, A. Peters, S. Velten, M.W., Phys. Rev. D **103**, 114506 (2021) [arXiv:2101.00723]]

– **Meson-meson operator** ( $BB$ ):

$$\mathcal{O}_{BB,\Gamma} = 2N_{BB}(\mathcal{C}\Gamma)_{AB}(\mathcal{C}\tilde{\Gamma})_{CD} \left( \bar{Q}_C^a(-\mathbf{r}/2)\psi_A^{(f)a}(-\mathbf{r}/2) \right) \left( \bar{Q}_D^b(+\mathbf{r}/2)\psi_B^{(f')b}(+\mathbf{r}/2) \right)$$

with  $\Gamma \in \{(1 + \gamma_0)\gamma_5, \gamma_5\}$  ( $\rightarrow (j_z, \mathcal{P}, \mathcal{P}_x) = (0, -, +)$ ).


– **Diquark-antidiquark operator** ( $Dd$ ):

$$\mathcal{O}_{Dd,\Gamma} = -N_{Dd}\epsilon^{abc} \left( \psi_A^{(f)b}(\mathbf{z})(\mathcal{C}\Gamma)_{AB}\psi_B^{(f')c}(\mathbf{z}) \right) \epsilon^{ade} \left( \bar{Q}_C^f(-\mathbf{r}/2)U^{fd}(-\mathbf{r}/2; \mathbf{z})(\mathcal{C}\tilde{\Gamma})_{CD}\bar{Q}_D^g(+\mathbf{r}/2)U^{ge}(+\mathbf{r}/2; \mathbf{z}) \right)$$

with  $\Gamma \in \{(1 + \gamma_0)\gamma_5, \gamma_5\}$  ( $\rightarrow (j_z, \mathcal{P}, \mathcal{P}_x) = (0, -, +)$ ).

- $\psi^{(f)}\psi^{(f')} = ud - du$  ( $\rightarrow I = 0$ ).
- $\tilde{\Gamma} = (1 + \gamma_0)\gamma_3$  (essentially irrelevant).
- Compute the  $4 \times 4$  correlation matrix

$$C_{jk}(t) = \langle \Omega | \mathcal{O}_j^\dagger(t) \mathcal{O}_k(0) | \Omega \rangle.$$







# Structure of the $\bar{b}\bar{b}qq$ tetraquark (2)

- Effective energies corresponding to diagonal elements of the correlation matrix,

$$V_j^{\text{eff}}(r, t) = -\frac{1}{a} \log \left( \frac{C_{jj}(t)}{C_{jj}(t-a)} \right) \quad (\text{no sum over } j).$$

- For large  $\bar{b}\bar{b}$  separations (right plot  $r \approx 0.79$  fm),  $BB$  effective energies reach plateaus at smaller  $t$  separations than  $Dd$  effective energies.  
 →  $BB$  dominates at large  $r$ ,  $Dd$  not important (energetically disfavored due to flux tube).
- For small  $\bar{b}\bar{b}$  separations (left plot  $r \approx 0.16$  fm),  $BB$  and  $Dd$  effective energies similar.  
 → More detailed investigation at small  $r$  necessary.




# Structure of the $\bar{b}\bar{b}qq$ tetraquark (3)

- Differences of effective energies corresponding to diagonal elements of the correlation matrix at small temporal separation  $t = 2a$  as functions of the  $\bar{b}\bar{b}$  separation  $r$ ,

$$V_j^{\text{eff}}(r, t = 2a) - V_k^{\text{eff}}(r, t = 2a).$$

- $BB$  versus  $Dd$  (left):  $Dd$  dominates for  $r \lesssim 3.15 a \approx 0.25 \text{ fm}$ , while  $BB$  dominates for  $r \gtrsim 3.15 a \approx 0.25 \text{ fm}$ .
- $BB$  operators (center):  $\Gamma = (1 + \gamma_0)\gamma_5$  leads to larger ground state overlap than  $\Gamma = \gamma_5$ . (Expected. Via a Fierz transformation one can show that  $\Gamma = (1 + \gamma_0)\gamma_5$  generates exclusively ground state mesons, while  $\gamma_5$  also generates parity excitations.)
- $Dd$  operators (right):  $\Gamma = (1 + \gamma_0)\gamma_5$  leads to larger ground state overlap than  $\Gamma = \gamma_5$ . (Interesting. In the literature mostly  $\gamma_5$  is discussed.)



# Structure of the $\bar{b}\bar{b}qq$ tetraquark (4)

- Optimize trial states

$$|\Phi_{b,d}\rangle = |b\rangle|\Phi_{BB,(1+\gamma_0)\gamma_5}\rangle + |d\rangle|\Phi_{Dd,(1+\gamma_0)\gamma_5}\rangle$$

( $|\Phi_j\rangle = \mathcal{O}_j|\Omega\rangle$ ) by minimizing effective energies


$$V_{b,d}^{\text{eff}}(r, t) = -\frac{1}{a} \log \left( \frac{C_{[b,d][b,d]}(t)}{C_{[b,d][b,d]}(t-a)} \right), \quad C_{[b,d][b,d]}(t) = \begin{pmatrix} b \\ d \end{pmatrix}_j^\dagger C_{jk}(t) \begin{pmatrix} b \\ d \end{pmatrix}_k.$$

with respect to  $b, d \in \mathbb{C}$ .

- Since norm and phase of  $b$  and  $d$  are irrelevant, consider relative weights of  $BB$  and  $Dd$ ,


$$w_{BB} = \frac{|b|^2}{|b|^2 + |d|^2}, \quad w_{Dd} = \frac{|d|^2}{|b|^2 + |d|^2} = 1 - w_{BB}.$$

- For fixed  $\bar{b}\bar{b}$  separation  $r$ ,  $w_{BB}$  and  $w_{Dd}$  depend only weakly on  $t$ .  
 $\rightarrow w_{BB}$  and  $w_{Dd}$  estimate the percentage of  $BB$  and of  $Dd$ .



# Structure of the $\bar{b}\bar{b}qq$ tetraquark (5)

- $w_{BB}$  and  $w_{Dd}$  as functions of the  $\bar{b}\bar{b}$  separation  $r$  (for two ensembles,  $a \approx 0.079$  fm and  $a \approx 0.063$  fm).
- $r \lesssim 0.2$  fm: Clear diquark-antidiquark dominance.
- $0.2 \text{ fm} \lesssim r \lesssim 0.3 \text{ fm}$ : Diquark-antidiquark dominance turns into meson-meson dominance.
- $0.5 \text{ fm} \lesssim r$ : Essentially a meson-meson system.



# Structure of the $\bar{b}\bar{b}qq$ tetraquark (6)

- Generalized eigenvalue problem (GEVP)

$$C_{jk}(t)v_k^{(n)}(t) = \lambda^{(n)}(t)C_{jk}(t_0)v_k^{(n)}(t_0) \quad , \quad n = 0, \dots, N-1$$

for  $t_0/a \geq 1$  and  $t/a > t_0/a$  with corresponding effective energies

$$V^{\text{eff},(n)}(r, t) = -\frac{1}{a} \log \left( \frac{\lambda^{(n)}(t)}{\lambda^{(n)}(t-a)} \right).$$

- **Eigenvector components**  $v_j^{(n)}(t)$  (which we always normalize according to  $\sum_j |v_j^{(n)}(t)|^2 = 1$ ) contain information about the relative importance of the operators. For large  $t$  and  $t_0$ ,

$$|n\rangle \approx \sum_j v_j^{(n)}(t) |\Phi_j\rangle,$$

where  $\approx$  denotes an approximate expansion of the **energy eigenstate**  $|n\rangle$  in terms of the trial states  $|\Phi_j\rangle$ .

# Structure of the $\bar{b}\bar{b}q q$ tetraquark (7)

- One can show: For  $t_0 = t - a$ , optimizing trial states by minimizing effective energies (as on previous slides) is equivalent to solving a GEVP, i.e.


$$(w_{BB}, w_{Dd}) = (|v_{BB,(1+\gamma_0)\gamma_5}^{(0)}|^2, |v_{Dd,(1+\gamma_0)\gamma_5}^{(0)}|^2)$$

(might offer another perspective on GEVP eigenvector components).

→ Results for  $w_{BB}$  and  $w_{Dd}$  can also be interpreted as GEVP results.

[P. Bicudo, A. Peters, S. Velten, M.W., Phys. Rev. D **103**, 114506 (2021) [arXiv:2101.00723]]

- In the literature typically small values for  $t_0$  are used, e.g.  $t_0/a = 1$  (instead of  $t_0 = t - a$  as used to obtain  $w_{BB}$  and  $w_{Dd}$  on previous slides).
- Similar results also for  $t_0/a = 1$ , when using a  $2 \times 2$  correlation matrix (left plot).
- Consistent results, when using a  $4 \times 4$  correlation matrix (right plot).



# Structure of the $\bar{b}\bar{b}qq$ tetraquark (8)

- Define the  $r$  dependent  $BB$  and  $Dd$  percentages,

$$p_{BB}(r) = w_{BB} \quad , \quad p_{Dd}(r) = w_{Dd}$$

and use the probability density of the  $\bar{b}\bar{b}$  separation

$$p_r(r) = 4\pi|R(r)|^2$$

obtained from the BO wave function  $R(r)/r$ , to estimate the total  $BB$  and  $Dd$  percentages of the  $\bar{b}\bar{b}ud$  tetraquark with quantum numbers  $I(J^P) = 0(1^+)$ :

$$\%BB = \int dr p_r(r)p_{BB}(r) \quad , \quad \%Dd = \int dr p_r(r)p_{Dd}(r) = 1 - \%BB.$$

- We find  $\%BB = 0.58$ ,  $\%Dd = 0.42$ .
- Using  $|v_{BB,(1+\gamma_0)\gamma_5}^{(0)}|^2$ ,  $|v_{Dd,(1+\gamma_0)\gamma_5}^{(0)}|^2$  instead of  $w_{BB}, w_{Dd}$  we find  $\%BB = 0.60$ ,  $\%Dd = 0.40$ .
- Results are in qualitative agreement with a GEVP result we obtained in a full lattice QCD computation, where the  $\bar{b}$  quarks are treated within NRQCD.

[L. Leskovec, S. Meinel, M. Pflaumer and M.W., Phys. Rev. D **100**, 014503 (2019) [arXiv:1904.04197]]  
[Talk by M. Pflaumer, today in this session at 6:45 a.m.]

# Summary

- The hadronically stable  $\bar{b}b\bar{u}d$  tetraquark with quantum numbers  $I(J^P) = 0(1^+)$  is neither exclusively a meson-meson system nor a diquark-antidiquark pair.
- $r \lesssim 0.2$  fm: Clear diquark-antidiquark dominance.
- $r \gtrsim 0.3$  fm: Clear meson-meson dominance.
- Total  $BB$  and  $Dd$  percentages:  $\%BB \approx 0.60$ ,  $\%Dd \approx 0.40$ .

