

Exercise sheet I

April 20 [correction: April 27]

Problem 1 [Zassenhaus formula]

Using the Zassenhaus formula $e^{A+B} = e^A e^B e^{Z_2} e^{Z_3} e^{Z_4} \dots$ with $Z_2 = \frac{1}{2}[B, A]$, $Z_3 = -\frac{1}{6}(2[B, [B, A]] + [A, [B, A]])$ and exponents $Z_n, n > 3$ containing higher powers of A and B , show the relation

$$e^{-i\epsilon H} = e^{-i\epsilon(H_0+V)} = e^{-i\epsilon V/2} e^{-i\epsilon H_0} e^{-i\epsilon V/2} + \mathcal{O}(\epsilon^3). \quad (1)$$

Problem 2 [Harmonic oscillator – partition function]

You are to obtain the partition function of the (quantum mechanical and classical) harmonic oscillator in three different ways:

- (i) Starting from the formula of the harmonic oscillator seen in the Lectures

$$\langle x, t | x, 0 \rangle = \sqrt{\frac{m\omega}{2\pi i \sin(\omega t)}} \exp\left(i \frac{x^2 m\omega}{\sin(\omega t)} (\cos(\omega t) - 1)\right) \quad (2)$$

go to Euclidean time ($t = -i\tau$) and calculate the partition function $Z = \int dx \langle x | e^{-\tau H} | x \rangle$.

- (ii) The oscillator spectrum is discrete. Make use of this fact and calculate $Z = \sum_n e^{-\tau E_n}$. Compare to (i).

- (iii) Now calculate the partition function of the classical oscillator using $Z = \int dx dp e^{-\tau(T+V)}$.

Compute the free energy $F = -1/\tau \log Z$ for the three cases and show that they agree for $\tau \ll 1$. Why?

Problem 3 [Functional derivative, chain rule]

Calculate the functional derivative, with respect to $f(x)$, of the functional $F[g(f)] = \int dx g(f(x))$ with $g(f) = \left(\frac{d}{dx} f(x)\right)^n$ using:

- (i) the definition of the (functional) chain rule $\frac{\delta F[g(\phi)]}{\delta \phi(y)} = \int ds \frac{\delta F[g(\phi)]}{\delta g(\phi(s))} \frac{\delta g(\phi(s))}{\delta \phi(y)}$;

- (ii) the chain rule of ordinary differential calculus and the equivalent definition $\frac{\delta F[\phi]}{\delta \phi(y)} = \left. \frac{d}{ds} F[\phi(x) + s\delta(x-y)] \right|_{s=0}$.